

➤ Bibliography

- [1] Aarjane, M., Slassi, S., Tazi, B., Maouloua, M., & Amine, A. (2019). Novel series of acridone-1,2,3-triazole derivatives: microwave-assisted synthesis, DFT study and antibacterial activities. *Journal of Chemical Sciences*, 131(8). <https://doi.org/10.1007/s12039-019-1653-2>
- [2] Abbas, I. M., Riyadh, S. M., Abdallah, M. A., & Gomha, S. M. (2006). A novel route to tetracyclic fused tetrazines and thiadiazines. *Journal of Heterocyclic Chemistry*, 43(4), 935–942. <https://doi.org/10.1002/jhet.5570430419>
- [3] Abd El-Aal, H. A. K., El-Khawaga, A. M. A. M., & Khalaf, A. A. (2014). Friedel-Crafts chemistry: Part 41. A new facile synthesis of indeno[1,2-c] pyrazoles, 2H-benzo[g]indazoles and benzo[6,7]cyclohepta[1,2-c]pyrazoles via Friedel-Crafts ring closures. *European Journal of Chemistry*, 5(2), 277–286. <https://doi.org/10.5155/eurjchem.5.2.277-286.976>
- [4] Abdel-Aziz, M., Abuo-Rahma, G. E.-D. A., & Hassan, A. A. (2009). Synthesis of novel pyrazole derivatives and evaluation of their antidepressant and anticonvulsant activities. *European Journal of Medicinal Chemistry*, 44(9), 3480–3487. <https://doi.org/10.1016/j.ejmech.2009.01.032>
- [5] Abdel-Aziz, S. A.-G., Ali, T. E.-S., El-Mahdy, K. M., & Abdel-Karim, S. M. (2011). Synthesis and antimicrobial activities of some novel bis-pyrazole derivatives containing a hydrophosphoryl unit. *European Journal of Chemistry*, 2(1), 25–35. <https://doi.org/10.5155/eurjchem.2.1.25-35.208>
- [6] Abdelrehim, E. M. (2021). Synthesis and Screening of New [1,3,4]Oxadiazole, [1,2,4]Triazole, and [1,2,4]Triazolo[4,3- b][1,2,4]triazole Derivatives as Potential Antitumor Agents on the Colon Carcinoma Cell Line (HCT-116). *ACS Omega*, 6(2), 1687–1696. <https://doi.org/10.1021/acsomega.0c05718>
- [7] Abdel-Wahab, B. F., Khidre, R. E., Farahat, A. A., & El-Ahl, A. S. (2012). ChemInform Abstract: 2-Chloroquinoline-3-carbaldehydes: Synthesis, Reactions and Applications. *ChemInform*, 43(30), 1–30. <https://doi.org/10.1002/chin.201230255>
- [8] Abdou, W. M., Khidre, R. E., & Kamel, A. A. (2012). Elaborating on Efficient Anti-Proliferation Agents of Cancer Cells and Anti-Inflammatory-Based N

- Bisphosphonic Acids. *Archiv Der Pharmazie*, 345(2), 123–136. <https://doi.org/10.1002/ardp.201100080>
- [9] Abdou, W. M., Khidre, R. E., & Shaddy, A. A. (2013). Synthesis of Tetrazoloquinoline-Based Mono- and Bisphosphonate Esters as Potent Anti-Inflammatory Agents. *Journal of Heterocyclic Chemistry*, 50(1), 33–41. <https://doi.org/10.1002/jhet.968>
- [10] Abdul Rahman, S. M., Bhatti, J. S., Thareja, S., & Monga, V. (2023). Current development of 1,2,3-triazole derived potential antimalarial scaffolds: Structure-activity relationship (SAR) and bioactive compounds. *European Journal of Medicinal Chemistry*, 259, 115699. <https://doi.org/10.1016/j.ejmech.2023.115699>
- [11] Aboul-Fadl, T., Mohammed, F. A.-Hamid., & Hassan, E. A.-S. (2003). Synthesis, antitubercular activity and pharmacokinetic studies of some schiff bases derived from 1- alkylisatin and isonicotinic acid hydrazide (inh). *Archives of Pharmacal Research*, 26(10), 778–784. <https://doi.org/10.1007/BF02980020>
- [12] Abramov, M. A., Ceulemans, E., Jackers, C., Van der Auweraer, M., & Dehaen, W. (2001). Reactions of 5-amino-1,2-azoles with aromatic and heterocyclic o-chloroaldehydes: [1+1] versus [2+1] cyclocondensation. *Tetrahedron*, 57(44), 9123–9129. [https://doi.org/10.1016/S0040-4020\(01\)00917-6](https://doi.org/10.1016/S0040-4020(01)00917-6)
- [13] Abu-Dief, A. M., & Mohamed, I. M. A. (2015). A review on versatile applications of transition metal complexes incorporating Schiff bases. *Journal of Basic and Applied Sciences*, 4(2), 119–133. <https://doi.org/10.1016/j.bjbas.2015.05.004>
- [14] Adagu, I. S. (2002). In vitro activity of nitazoxanide and related compounds against isolates of *Giardia intestinalis*, *Entamoeba histolytica* and *Trichomonas vaginalis*. *Journal of Antimicrobial Chemotherapy*, 49(1), 103–111. <https://doi.org/10.1093/jac/49.1.103>
- [15] Aday, H. A. (2013). Synthesis and Characterization of the Triazole Derived from Thiosemicarbazide, 4-Amino-5-Phenyl-4H-1,2,4-Triazole-3-Thiol and Their Copper(II) and Nickel(II) Complexes. *Engineering and Technology Journal*, 31(2). www.pdfactory.com
- [16] Afzal, O., Kumar, S., Haider, M. R., Ali, M. R., Kumar, R., Jaggi, M., & Bawa, S. (2015). A review on anticancer potential of bioactive heterocycle quinoline. *European*

- Journal of Medicinal Chemistry*, 97, 871–910.
<https://doi.org/10.1016/j.ejmech.2014.07.044>
- [17] Akhramez, S., Hafid, A., Khouili, M., Saadi, M., Ammari, L. El, & Ketatni, E. M. (2019). Synthesis, crystal structure and Hirshfeld surface analysis of 2-chloro-3-[(E)-(2-phenylhydrazinylidene)methyl]quinoline. *Acta Crystallographica Section E: Crystallographic Communications*, 75, 964–968.
<https://doi.org/10.1107/S2056989019007692>
- [18] Alhaider, A. A., Abdelkader, M. A., & Lien, E. J. (1986). Design, Synthesis, and Pharmacological Activities of 2-Substituted 4-Phenyl- quinolines as Potential Antidepressant Drugs. *Chemischer Informations Dienst*, 17(5).
<https://doi.org/10.1002/chin.198605200>
- [19] Ali, S. M. M., Azad, M. A. K., Jesmin, M., Ahsan, S., Rahman, M. M., Khanam, J. A., Islam, M. N., & Shahriar, S. M. S. (2012). In vivo anticancer activity of vanillin semicarbazone. *Asian Pacific Journal of Tropical Biomedicine*, 2(6), 438–442.
[https://doi.org/10.1016/S2221-1691\(12\)60072-0](https://doi.org/10.1016/S2221-1691(12)60072-0)
- [20] Al-Messri, A. (2009). Synthesis of Some New 1,2,4-Triazoles Derived from 2-Mercaptobenzimidazole. *Um-Salama Science Journal*, 6(1).
- [21] Al-Sha'alan, N. (2007). Antimicrobial Activity and Spectral, Magnetic and Thermal Studies of Some Transition Metal Complexes of a Schiff Base Hydrazone Containing a Quinoline Moiety. *Molecules*, 12(5), 1080–1091. <https://doi.org/10.3390/12051080>
- [22] Al-Soud, Y. A., Al-Masoudi, N. A., & Ferwanah, A. E. R. S. (2003). Synthesis and properties of new substituted 1,2,4-triazoles: Potential antitumor agents. *Bioorganic and Medicinal Chemistry*, 11(8), 1701–1708. [https://doi.org/10.1016/S0968-0896\(03\)00043-9](https://doi.org/10.1016/S0968-0896(03)00043-9)
- [23] Altimari, J. M., Hockey, S. C., Boshoff, H. I., Sajid, A., & Henderson, L. C. (n.d.). *Supporting Information Novel 1,4-Substituted-1,2,3-Triazoles as Antitubercular Agents*.
- [24] Aly, E.-S. A. and E.-B. M. A. and B. M. A. (2004). A convenient synthesis of some novel pyrazole derivatives. *Indian Journal of Chemistry. Sect. B: Organic Chemistry, Including Medical Chemistry*, 43(6), 1355–1359.

- [25] Amir, M., Javed, S. A., & Kumar, H. (2007). Synthesis of Some 1,3,4-Oxadiazole Derivatives as Potential Antiinflammatory Agents. *ChemInform*, 38(39). <https://doi.org/10.1002/chin.200739109>
- [26] Anandarajagopal, K., Anbu, J., Sunilson, J., Illavarasu, A., Thangavelpandian, N., & Kalirajan, R. (2010). Antiepileptic and Antimicrobial Activities of Novel 1-(unsubstituted/substituted)-3,5-dimethyl-1H-pyrazole Derivatives. *International Journal of ChemTech Research*, 2(1), 45–49.
- [27] Angajala, K. K., Vianala, S., Macha, R., Raghavender, M., Thupurani, M. K., & Pathi, P. J. (2016). Synthesis, anti-inflammatory, bactericidal activities and docking studies of novel 1,2,3-triazoles derived from ibuprofen using click chemistry. *SpringerPlus*, 5(1), 423. <https://doi.org/10.1186/s40064-016-2052-5>
- [28] Ansari, A., Ali, A., Asif, M., & Shamsuzzaman, S. (2018). Microwave-assisted MgO NP catalyzed one-pot multicomponent synthesis of polysubstituted steroidal pyridines. *New Journal of Chemistry*, 42(1), 184–197. <https://doi.org/10.1039/C7NJ03742B>
- [29] Arthur Michael, von. (1893). Ueber die Einwirkung von Diazobenzolimid anf Acetylendicarbonsauremethylester. *Annalen Der Chemie*, 273, 94–95.
- [30] As, T., Sal, S., & Ha, T. (2016). Evaluation of Novel 2-Chloro Quinoline-3-Carbaldehyde derivatives. *International Journal of Chemical Studies*, 4(6), 95–98.
- [31] Ashry, E. S. H. El, Elshatanofy, M. M., Badawy, M. E. I., Kandeel, K. M., Elhady, O. M., & Abdel-Sayed, M. A. (2020). Synthesis and Evaluation of Antioxidant, Antibacterial, and Target Protein-Molecular Docking of Novel 5-Phenyl-2,4-dihydro-3H-1,2,4-triazole Derivatives Hybridized with 1,2,3-Triazole via the Flexible SCH2-Bonding. *Russian Journal of General Chemistry*, 90(12), 2419–2434. <https://doi.org/10.1134/S1070363220120300>
- [32] Askar, F. W., Hassan, H. A., & Jinzeel, N. A. (2013). Synthesis of Some Heterocyclic Compounds Derived from 2-Mercapto Benzoxazole. *Baghdad Science Journal*, 10(3), 766–778.
- [33] Athar Abbasi, M., Akhtar, A., Nafeesa, K., Zahra Siddiqui, S., Mohammed Khan, K., Ashraf, M., & Abida Ejaz, S. (2013). Synthesis, structural characterization and enzyme inhibition studies on 5-(2-nitrostyryl)-1,3,4-oxadiazole-2-thiol derivatives. *Journal of Chilean Chemical Society*, 58(4), 2186–2191.

- [34] Avaji, P. G., Vinod Kumar, C. H., Patil, S. A., Shivananda, K. N., & Nagaraju, C. (2009). Synthesis, spectral characterization, in-vitro microbiological evaluation and cytotoxic activities of novel macrocyclic bis hydrazone. *European Journal of Medicinal Chemistry*, 44(9), 3552–3559. <https://doi.org/10.1016/j.ejmech.2009.03.032>
- [35] Bagdi, A. K., Santra, S., Monir, K., & Hajra, A. (2015). Synthesis of imidazo[1,2-a]pyridines: a decade update. *Chemical Communications*, 51(9), 1555–1575. <https://doi.org/10.1039/C4CC08495K>
- [36] Bakht, M. A., Yar, M. S., Abdel-Hamid, S. G., Al Qasoumi, S. I., & Samad, A. (2010). Molecular properties prediction, synthesis and antimicrobial activity of some newer oxadiazole derivatives. *European Journal of Medicinal Chemistry*, 45(12), 5862–5869. <https://doi.org/10.1016/j.ejmech.2010.07.069>
- [37] Balabin, R. M. (2009). Tautomeric equilibrium and hydrogen shifts in tetrazole and triazoles: Focal-point analysis and ab initio limit. *Journal of Chemical Physics*, 131(15). <https://doi.org/10.1063/1.3249968>
- [38] Balakrishna Prabhu, K., Saidutta, M. B., Isloor, A. M., & Hebbar, R. (2017). Improvement in performance of polysulfone membranes through the incorporation of chitosan-(3-phenyl-1h-pyrazole-4-carbaldehyde). *Cogent Engineering*, 4(1). <https://doi.org/10.1080/23311916.2017.1403005>
- [39] Baraldi, P. G., Fruttarolo, F., Tabrizi, M. A., Romagnoli, R., & Preti, D. (2007). Novel 8-heterocyclyl xanthine derivatives in drug development - An update. *Expert Opinion on Drug Discovery*, 2(9), 1161–1183. <https://doi.org/10.1517/17460441.2.9.1161>
- [40] Barnes, P. J. (2013). Theophylline. *American Journal of Respiratory and Critical Care Medicine*, 188(8), 901–906. <https://doi.org/10.1164/rccm.201302-0388PP>
- [41] Barus, M., Rotar, D., Bratenko, M., Panasenko, N., Zvarych, V., Stasevych, M., & Vovk, M. (2021). Synthesis and evaluation of antimicrobial activities of new functional derivatives of 3-[5-(4-nitrophenyl)-2-furyl]-4-pyrazole-carbaldehydes. In *Biointerface Research in Applied Chemistry* (Vol. 11, Issue 4, pp. 12159–12169). AMG Transcend Association. <https://doi.org/10.33263/BRIAC114.1215912169>

- [42] Bawa, S., & Kumar, S. (2009). Synthesis of Schiff's bases of 8-methyl-tetrazolo[1,5-a]quinoline as potential anti-inflammatory and antimicrobial agents. *Indian Journal of Chemistry*, 48, 142–145.
- [43] Baziard-Mouysset, G., Rached, A., Younes, S., Tournaire, C., Stigliani, J., Payard, M., Yavo, J., & Advenier, C. (1995). Synthesis and in vitro bronchospasmolytic activity of 8-aryl, heteroaryl or arylalkyl theophyllines. *European Journal of Medicinal Chemistry*, 30(3), 253–260. [https://doi.org/10.1016/0223-5234\(96\)88233-X](https://doi.org/10.1016/0223-5234(96)88233-X)
- [44] Bekircan, O., Menteşe, E., Ülker, S., & Kucuk, C. (2014). Synthesis of some new 1,2,4-triazole derivatives starting from 3-(4-Chlorophenyl)-5-(4-methoxybenzyl)-4H-1,2,4-triazol with anti-lipase and anti-urease activities. *Archiv Der Pharmazie*, 347(6), 387–397. <https://doi.org/10.1002/ardp.201300344>
- [45] Belkheira, M., El Abed, D., Pons, J., & Bressy, C. (2011). Organocatalytic Synthesis of 1,2,3-Triazoles from Unactivated Ketones and Arylazides. *Chemistry – A European Journal*, 17(46), 12917–12921. <https://doi.org/10.1002/chem.201102046>
- [46] Berk, B., Akgün, H., Erol, K., Sirmagül, B., Gao, Z. G., & Jacobson, K. A. (2005). New 8-substituted xanthine derivatives as potent bronchodilators. *Farmaco*, 60(11–12), 974–980. <https://doi.org/10.1016/j.farmac.2005.08.011>
- [47] Beyzaei, H., Ghanbari Kudeyani, M., Samareh Delarami, H., & Aryan, R. (2020). Synthesis, antimicrobial and antioxidant evaluation, and molecular docking study of 4,5-disubstituted 1,2,4-triazole-3-thiones. *Journal of Molecular Structure*, 1215. <https://doi.org/10.1016/j.molstruc.2020.128273>
- [48] Beyzaei, H., Khosravi, Z., Aryan, R., & Ghasemi, B. (2019). A green one-pot synthesis of 3(5)-substituted 1,2,4-triazol-5(3)-amines as potential antimicrobial agents. *Journal of the Iranian Chemical Society*, 16(12), 2565–2573. <https://doi.org/10.1007/s13738-019-01714-2>
- [49] Bhagat, S. B., & Telvekar, V. N. (2017a). NBS mediated protocol for the synthesis of N -bridged fused heterocycles in water. *Tetrahedron Letters*, 58(37), 3662–3666. <https://doi.org/10.1016/j.tetlet.2017.08.017>
- [50] Bhagat, S. B., & Telvekar, V. N. (2017b). NBS mediated protocol for the synthesis of N-bridged fused heterocycles in water. *Tetrahedron Letters*, 58(37), 3662–3666. <https://doi.org/10.1016/j.tetlet.2017.08.017>

- [51] Bhardwaj, V., Noolvi, M. N., Jalhan, S., & Patel, H. M. (2016). Synthesis, and antimicrobial evaluation of new pyridine imidazo [2,1b]-1,3,4-thiadiazole derivatives. *Journal of Saudi Chemical Society*, 20, S406–S410. <https://doi.org/10.1016/j.jscs.2012.12.007>
- [52] Bhovi, V. K., Bodke, Y. D., Biradar, S., Swamy, B. E. K., & Umesh, S. (2009). A facile synthesis of bromo-substituted benzofuran containing thiazolidinone nucleus bridged with quinoline derivatives: Potent analgesic and antimicrobial agents. *Phosphorus, Sulfur and Silicon and the Related Elements*, 185(1), 110–116. <https://doi.org/10.1080/10426500902717317>
- [53] Bhutani, R., Pathak, D. P., Kapoor, G., Husain, A., & Iqbal, M. A. (2019). Novel hybrids of benzothiazole-1,3,4-oxadiazole-4-thiazolidinone: Synthesis, in silico ADME study, molecular docking and in vivo anti-diabetic assessment. *Bioorganic Chemistry*, 83, 6–19. <https://doi.org/10.1016/j.bioorg.2018.10.025>
- [54] Biscussi, B., & Murray, A. P. (2022). New Caffeine Derivatives as Multitarget Agents for the Therapy of Alzheimer's Disease †. *Chem. Proc*, 2022. <https://doi.org/10.3390/xxxxx>
- [55] Boceiri, N., Benabdallah, T., Hadj Youcef, M., & Reffas, H. (2016). Synthesis and Characterization of a Novel Series of Amphiphilic Mercapto-1,2,4-Triazole Schiff Base Ligands: Investigation of their Behavior in Hydro-Organic Solutions. *Journal of Surfactants and Detergents*, 19(3), 583–597. <https://doi.org/10.1007/s11743-016-1811-1>
- [56] Bolakatti, G., Ramesh, D. K., Gopala Krishna, B., Mruthyunjaya, J. H., & Katagi, M. S. (2021). Synthesis and Antimicrobial Evaluation of Newly Sythesized Triazolothiadiazole Analogs. *International Journal of Drug Design and Discovery*, 4(2), 1050–1055. <https://www.researchgate.net/publication/353120984>
- [57] Bondock, S., Gieman, H., & El-Shafei, A. (2016). Selective synthesis, structural studies and antitumor evaluation of some novel unsymmetrical 1-hetaryl-4-(2-chloroquinolin-3-yl)azines. *Journal of Saudi Chemical Society*, 20(6), 695–702. <https://doi.org/10.1016/j.jscs.2015.01.005>
- [58] Borowiecki, P., Paprocki, D., Dudzik, A., & Plenkiewicz, J. (2016). Chemoenzymatic Synthesis of Proxiphylline Enantiomers. *Journal of Organic Chemistry*, 81(2), 380–395. <https://doi.org/10.1021/acs.joc.5b01840>

- [59] Borowiecki, P., Rudzka, A., Reiter, T., & Kroutil, W. (2022). Biocatalytic hydrogen-transfer to access enantiomerically pure proxyphylline, xanthinol, and diprophylline. *Bioorganic Chemistry*, 127. <https://doi.org/10.1016/j.bioorg.2022.105967>
- [60] Boukhssas, S., Aouine, Y., Faraj, H., Alami, A., El Hallaoui, A., & Bekkari, H. (2017). Synthesis, Characterization, and Antibacterial Activity of Diethyl 1-((4-Methyl-2-phenyl-4,5-dihydrooxazol-4-yl)methyl)-1H-1,2,3-triazole-4,5-dicarboxylate. *Journal of Chemistry*, 2017. <https://doi.org/10.1155/2017/4238360>
- [61] Bulbule, V. J., Deshpande, V. H., Velu, S., Sudalai, A., Sivasankar, S., & Sathe, V. T. (1999). Heterogeneous Henry reaction of aldehydes: Diastereoselective synthesis of nitroalcohol derivatives over Mg-Al hydrotalcites. *Tetrahedron*, 55(30), 9325–9332. [https://doi.org/10.1016/S0040-4020\(99\)00494-9](https://doi.org/10.1016/S0040-4020(99)00494-9)
- [62] Bulut, N., Kocyigit, U. M., Gecibesler, I. H., Dastan, T., Karci, H., Taslimi, P., Durna Dastan, S., Gulcin, I., & Cetin, A. (2018). Synthesis of some novel pyridine compounds containing bis-1,2,4-triazole/thiosemicarbazide moiety and investigation of their antioxidant properties, carbonic anhydrase, and acetylcholinesterase enzymes inhibition profiles. *Journal of Biochemical and Molecular Toxicology*, 32(1). <https://doi.org/10.1002/jbt.22006>
- [63] Byrnes, J. J., Miller, L. G., Greenblatt, D. J., & Shader, R. I. (1993). Chronic benzodiazepine administration. *Psychopharmacology*, 111(1), 91–95. <https://doi.org/10.1007/BF02257412>
- [64] Campbell, S. F., Hardstone, J. D., & Palmer, M. J. (1988). ChemInform Abstract: 2,4-Diamino-6,7-dimethoxyquinoline Derivatives as α 1-Adrenoceptor Antagonists and Antihypertensive Agents. *ChemInform*, 19(38). <https://doi.org/10.1002/chin.198838225>
- [65] Cao, H. T., & Gree, R. (2009). DEAD-(cat)ZnBr₂ an efficient system for the oxidation of alcohols to carbonyl compounds. *Tetrahedron Letters*, 50(13), 1493–1494. <https://doi.org/10.1016/j.tetlet.2009.01.080>
- [66] Cao, H. T., & Grée, R. (2009). DEAD-(cat)ZnBr₂ an efficient system for the oxidation of alcohols to carbonyl compounds. *Tetrahedron Letters*, 50(13), 1493–1494. <https://doi.org/10.1016/j.tetlet.2009.01.080>

- [67] Carmona, A. T., Carrion-Jimenez, S., Pingitore, V., Moreno-Clavijo, E., Robina, I., & Moreno-Vargas, A. J. (2018). Harnessing pyrrolidine iminosugars into dimeric structures for the rapid discovery of divalent glycosidase inhibitors. *European Journal of Medicinal Chemistry*, 151, 765–776. <https://doi.org/10.1016/j.ejmech.2018.04.008>
- [68] Chambhare, R. (2003). Synthesis and preliminary evaluation of some N-[5-(2-furanyl)-2-methyl-4-oxo-4H-thieno[2,3-d]pyrimidin-3-yl]-carboxamide and 3-substituted-5-(2-furanyl)-2-methyl-3H-thieno[2,3-d]pyrimidin-4-ones as antimicrobial agents. *European Journal of Medicinal Chemistry*, 38(1), 89–100. [https://doi.org/10.1016/S0223-5234\(02\)01442-3](https://doi.org/10.1016/S0223-5234(02)01442-3)
- [69] Chand, M., Kaushik, R., & Chand Jain, S. (2018). Synthesis and antimicrobial and antioxidant activities of hybrid molecules containing benzotriazole and 1,2,4-triazole. *Turkish Journal of Chemistry*, 42(6), 1663–1677. <https://doi.org/10.3906/kim-1803-61>
- [70] Chandramouli, C. , S. M. R. , N. T. B. , B. B. , & U. R. H. (2012). Synthesis and biological screening of certain new triazole Schiff bases and their derivatives bearing substituted benzothiazole moiety. *J. Chem. Pharm. Res*, 2(4), 1151–1159.
- [71] Chandrashekharappa, S. (2013). *Solvent-Free Solid Phase Syntheses of 2-Chloroquinoline-3-carbaldehyde Phenyl Hydrazones and their DNA Cleavage Studies*. www.joac.info
- [72] Charris, J., Lobo, G., Camacho, J., Ferrer, R., Barazarte, A., Dominguez, J., Gamboa, N., Rodrigues, J., & Angel, J. (2007). Synthesis and Antimalarial Activity of (E) 2-(2-Chloro-3-Quinolinylmethylidene)-5,7-Dimethoxyindanones. *Letters in Drug Design & Discovery*, 4(1), 49–54. <https://doi.org/10.2174/157018007778992865>
- [73] Chaubey, A. K. , & P. S. N. (2012). Synthesis & anticonvulsant activity (Chemo Shock) of Schiff and Mannich bases of Isatin derivatives with 2-Amino pyridine (mechanism of action). *International Journal of PharmTech Research*, 4(4), 590–598.
- [74] Chen, D., Li, J., Zhao, Y., & Wu, Y. (2022). Human Exposure of Fipronil Insecticide and the Associated Health Risk. *Journal of Agricultural and Food Chemistry*, 70(1), 63–71. <https://doi.org/10.1021/acs.jafc.1c05694>
- [75] Chen, Y.-L., Chen, I.-L., Lu, C.-M., Tzeng, C.-C., Tsao, L.-T., & Wang, J.-P. (2004). Synthesis and anti-inflammatory evaluation of 4-anilino-furo[2,3- b]quinoline and 4-

- phenoxyfuro[2,3- b]quinoline derivatives. Part 3. *Bioorganic & Medicinal Chemistry*, 12(2), 387–392. <https://doi.org/10.1016/j.bmc.2003.10.051>
- [76] Chinnasamy, R., Sundararajan, R., & Govindaraj, S. (2010). Synthesis, characterization, and analgesic activity of novel schiff base of isatin derivatives. *Journal of Advanced Pharmaceutical Technology & Research*, 1(3), 342. <https://doi.org/10.4103/0110-5558.72428>
- [77] Chitikina, S. S., Buddiga, P., Deb, P. K., Mailavaram, R. P., Venugopala, K. N., Nair, A. B., Al-Jaidi, B., & Kar, S. (2020). Synthesis and anthelmintic activity of some novel (E)-2-methyl/propyl-4-(2-(substitutedbenzylidene)hydrazinyl)-5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidines. *Medicinal Chemistry Research*, 29(9), 1600–1610. <https://doi.org/10.1007/s00044-020-02586-5>
- [78] Chlon-Rzepa, G., Slusarczyk, M., Jankowska, A., Gawalska, A., Bucki, A., Kolaczowski, M., Swierczek, A., Pocięcha, K., Wyska, E., Zygmunt, M., Kazek, G., Sałat, K., & Pawłowski, M. (2018). Novel amide derivatives of 1,3-dimethyl-2,6-dioxopurin-7-yl-alkylcarboxylic acids as multifunctional TRPA1 antagonists and PDE4/7 inhibitors: A new approach for the treatment of pain. *European Journal of Medicinal Chemistry*, 158, 517–533. <https://doi.org/10.1016/j.ejmech.2018.09.021>
- [79] Choudhare, T. S., Wagare, D. S., Shirsath, S. E., & Netankar, P. D. (2021). H 3 PO 4 catalyzed one-pot synthesis of 1,3-diphenyl-1H-pyrazole-4-carbaldehyde to novel 1,3-diphenyl-1H-pyrazole-4-carbonitrile. *Indian Academy of Sciences*, 69(133). <https://doi.org/10.1007/s12039-021>
- [80] Chunavala, K. C., Joshi, G., Suresh, E., & Adimurthy, S. (2011). Thermal and microwave-assisted rapid syntheses of substituted imidazo[1,2-a]pyridines under solvent- and catalyst-free conditions. *Thieme Verlag Stuttgart*, 2011(4), 635–641. <https://doi.org/10.1055/s-0030-1258405>
- [81] Cleary, J. D., Taylor, J. W., & Chapman, S. W. (1992). Itraconazole in Antifungal Therapy. *Annals of Pharmacotherapy*, 26(4), 502–509. <https://doi.org/10.1177/106002809202600411>
- [82] Corsano, S., Scapicchi, R., & Strappaghetti, G. (1994). Bronchodilator Activity of Theophylline Derivatives Substituted at the 7-Position. *Archiv Der Pharmazie*, 327(10), 631–635. <https://doi.org/10.1002/ardp.19943271006>

- [83] Cui, P., Li, X., Zhu, M., Wang, B., Liu, J., & Chen, H. (2017). Design, synthesis and antimicrobial activities of thiouracil derivatives containing triazolo-thiadiazole as SecA inhibitors. *European Journal of Medicinal Chemistry*, 127, 159–165. <https://doi.org/10.1016/j.ejmech.2016.12.053>
- [84] Cui, T., Zhang, X., Lin, J., Zhu, Z., Liu, P., & Sun, P. (2021). Electrochemical Oxidative C-H Thiocyanation or Selenocyanation of Imidazopyridines and Arenes. *Synlett*, 32(3), 267–272. <https://doi.org/10.1055/a-1299-3009>
- [85] Curioni, C., & Andre, C. (2006). Rimonabant for overweight or obesity. *Cochrane Database of Systematic Reviews*, 2010(1). <https://doi.org/10.1002/14651858.CD006162.pub2>
- [86] Cvietusa, P., Mascali, J. J., Negri, J., & Borish, L. (1996). Anti-inflammatory Effects of Theophylline: Modulation of Cytokine Production. *Annals of Allergy, Asthma & Immunology*, 77(1), 34–38. [https://doi.org/10.1016/S1081-1206\(10\)63476-X](https://doi.org/10.1016/S1081-1206(10)63476-X)
- [87] Cziaky, Z., Korodi, F., Frank, L., & Czink, I. (1996). Synthesis and antimycotic activity of new 2-chloro-3-(2-nitro)ethyl and (2-nitro)vinylquinolines. *Heterocyclic Communications*, 2(1). <https://doi.org/10.1515/HC.1996.2.1.63>
- [88] Da Silva, C. M., Da Silva, D. L., Modolo, L. V., Alves, R. B., De Resende, M. A., Martins, C. V. B., & De Fátima, Â. (2011). Schiff bases: A short review of their antimicrobial activities. *Journal of Advanced Research*, 2(1), 1–8. <https://doi.org/10.1016/j.jare.2010.05.004>
- [89] De, S. S., Khambete, M. P., & Degani, M. S. (2019). Oxadiazole scaffolds in anti-tuberculosis drug discovery. *Bioorganic and Medicinal Chemistry Letters*, 29(16), 1999–2007. <https://doi.org/10.1016/j.bmcl.2019.06.054>
- [90] Deb, P. K., Mailavaram, R., Chandrasekaran, B., Kaki, V. R., Kaur, R., Kachler, S., Klotz, K., & Akkinapally, R. R. (2018). Synthesis, adenosine receptor binding and molecular modelling studies of novel thieno[2,3-*d*]pyrimidine derivatives. *Chemical Biology & Drug Design*, 91(4), 962–969. <https://doi.org/10.1111/cbdd.13155>
- [91] Deep, A., Bhatia, R., Kaur, R., Kumar, S., Jain, U., Singh, H., Batra, S., Kaushik, D., & Deb, P. (2016). Imidazo[1,2-*a*]pyridine Scaffold as Prospective Therapeutic Agents. *Current Topics in Medicinal Chemistry*, 17(2), 238–250. <https://doi.org/10.2174/1568026616666160530153233>

- [92] Desai, N. C., & Dodiya, A. M. (2016). Conventional and microwave techniques for the synthesis and antimicrobial studies of novel 1-[2-(2-chloro-6-methyl(3-quinolyl))-5-(4-nitrophenyl)-(1,3,4-oxadiazolin-3-yl)]-3-(aryl)prop-2-en-1-ones. *Arabian Journal of Chemistry*, 9, S379–S387. <https://doi.org/10.1016/j.arabjc.2011.05.004>
- [93] Dheer, D., Singh, V., & Shankar, R. (2017). Medicinal attributes of 1,2,3-triazoles: Current developments. *Bioorganic Chemistry*, 71, 30–54. <https://doi.org/10.1016/j.bioorg.2017.01.010>
- [94] Dianov, V. M., & Bulgakov, A. K. (2006). Synthesis and antimicrobial activity of 3-methyl-substituted 6,8-dimethylthiazolo-[2,3-f]xanthines. *Pharmaceutical Chemistry Journal*, 40(10), 551–553. <https://doi.org/10.1007/s11094-006-0191-7>
- [95] Dubey, P. K., Rao, S. S., & Reddy, P. V. P. (2003). Synthesis of some novel 2-(2-chloro-3-quinolyl)-5-phenyl-2,3-dihydro-1,3,4-oxadiazoles. *Heterocyclic Communications*, 9(4), 411–414.
- [96] Dyminska, L. (2015). Imidazopyridines as a source of biological activity and their pharmacological potentials—Infrared and Raman spectroscopic evidence of their content in pharmaceuticals and plant materials. *Bioorganic & Medicinal Chemistry*, 23(18), 6087–6099. <https://doi.org/10.1016/j.bmc.2015.07.045>
- [97] El Abbouchi, A., Koubachi, J., El Brahmi, N., & Kazzouli, S. (2019). Direct arylation and Suzuki-Miyaura coupling of imidazo [1,2-a]pyridines catalyzed by (SIPr)Pd(allyl)Cl complex under microwave-irradiation. *Mediterranean Journal of Chemistry*, 9(5), 347–354. <https://doi.org/10.13171/mjc1911271124sek>
- [98] El Shehry, M. F., Abbas, S. Y., Farrag, A. M., Fouad, S. A., & Ammar, Y. A. (2020). Synthesis and biological evaluation of 3-(2,4-dichlorophenoxyethyl)-1-phenyl-1H-pyrazole derivatives as potential antitumor agents. *Journal of the Iranian Chemical Society*, 17(10), 2567–2575. <https://doi.org/10.1007/s13738-020-01951-w>
- [99] El-Emam, A. A., Alrashood, K. A., Al-Omar, M. A., & Al-Tamimi, A. M. S. (2012). Synthesis and antimicrobial activity of N'-heteroarylidene-1-adamantylcarbohydrazides and (±)-2-(1-adamantyl)-4-acetyl-5-[5-(4-substituted phenyl-3-isoxazolyl)]-1,3,4-oxadiazolines. *Molecules*, 17(3), 3475–3483. <https://doi.org/10.3390/molecules17033475>
- [100] El-Feky, S. A. H., Abd El-Samii, Z. K., Osman, N. A., Lashine, J., Kamel, M. A., & Thabet, H. Kh. (2015). Synthesis, molecular docking and anti-inflammatory screening

- of novel quinoline incorporated pyrazole derivatives using the Pfitzinger reaction II. *Bioorganic Chemistry*, 58, 104–116. <https://doi.org/10.1016/j.bioorg.2014.12.003>
- [101] Eliot, A. C., & Kirsch, J. F. (2004). Pyridoxal Phosphate Enzymes: Mechanistic, Structural, and Evolutionary Considerations. *Annual Review of Biochemistry*, 73(1), 383–415. <https://doi.org/10.1146/annurev.biochem.73.011303.074021>
- [102] Elsayed Khidre, R. and F. A.-W. B. and A.-R. B. F. (2011). New quinoline-based compounds for analgesic and anti-inflammatory evaluation. *Letters in Drug Design & Discovery*, 8(7), 640–648.
- [103] Espinoza-Vázquez, A., Rodríguez-Gómez, F. J., Martínez-Cruz, I. K., Ángeles-Beltrán, D., Negrón-Silva, G. E., Palomar-Pardavé, M., Romero, L. L., Pérez-Martínez, D., & Navarrete-López, A. M. (2019). Adsorption and corrosion inhibition behaviour of new theophylline-triazole-based derivatives for steel in acidic medium. *Royal Society Open Science*, 6(3). <https://doi.org/10.1098/rsos.181738>
- [104] Farghally, A. M. , H. N. S. , H. A. A. , & E.-S. O. A. (1985). Synthesis of substituted quinoline-3-carbaldehyde (2, 3-dihydrothiazol-2-ylidene) hydrazones of potential antimicrobial activity. *Journal de Pharmacie de Belgique*, 7(40), 366–372.
- [105] Farghaly, A. M., Habib, N. S., Khalil, M. A., El-Sayed, O. A., & Bistawroos, A. E. (1990). Synthesis of Novel 2-Substituted Quinoline Derivatives: Antimicrobial, Inotropic, and Chronotropic Activities. *Archiv Der Pharmazie*, 323(4), 247–251. <https://doi.org/10.1002/ardp.19903230414>
- [106] Fieger, S. M., & Wong, B. J. (2010). Adenosine receptor inhibition with theophylline attenuates the skin blood flow response to local heating in humans. *Experimental Physiology*, 95(9), 946–954. <https://doi.org/10.1113/expphysiol.2010.053538>
- [107] Fjellaksel Rune; Riss Patrick J.; Hansen Jørn H., R. S. (2018). Copper-Mediated Late-Stage Iodination and 123I-Labeling of Triazole-benzimidazole Bioactives. *Synlett*, 29(11), 1491–1495. <https://doi.org/10.1055/s-0036-1591985>
- [108] Friedrich, G. (2002). Determination of lonazolac and its hydroxy and O-sulfated metabolites by on-line sample preparation liquid chromatography with fluorescence detection. *Journal of Chromatography B*, 766(2), 295–305. [https://doi.org/10.1016/S0378-4347\(01\)00514-X](https://doi.org/10.1016/S0378-4347(01)00514-X)

- [109] Fu, H.-B., & Yao, J.-N. (2001). Size Effects on the Optical Properties of Organic Nanoparticles. *Journal of the American Chemical Society*, 123(7), 1434–1439. <https://doi.org/10.1021/ja0026298>
- [110] Gaidhane, M. K., Lemdeo, S., Mahavidhyalaya, P., Kuhl, M., Ghatole, A. M., & Lanjewar, K. R. (2013). Novel synthesis and antimicrobial activity of novel Schiff base derived quinolin and their β -lactum derivatives. <https://www.researchgate.net/publication/260419281>
- [111] Galgiani, J. N. (1990). Fluconazole, a New Antifungal Agent. *Annals of Internal Medicine*, 113(3), 177. <https://doi.org/10.7326/0003-4819-113-3-177>
- [112] Gamage, S. A., Spicer, J. A., Atwell, G. J., Finlay, G. J., Baguley, B. C., & Denny, W. A. (1999). Structure–Activity Relationships for Substituted Bis(acridine-4-carboxamides): A New Class of Anticancer Agents. *Journal of Medicinal Chemistry*, 42(13), 2383–2393. <https://doi.org/10.1021/jm980687m>
- [113] Gao, F., Wang, T., Xiao, J., & Huang, G. (2019). Antibacterial activity study of 1,2,4-triazole derivatives. *European Journal of Medicinal Chemistry*, 173, 274–281. <https://doi.org/10.1016/j.ejmech.2019.04.043>
- [114] Gao, Y. C., Huang, Z. H., Zhang, Z. S., Xie, J. X., Cui, Z. N., & Tang, R. Y. (2020). Sulfite-Promoted C-H Fluoroalkyl Sulfuration of Imidazoheterocycles with Bromofluoroacetate and Elemental Sulfur. *Synthesis (Germany)*, 52(17), 2541–2550. <https://doi.org/10.1055/s-0040-1707120>
- [115] Gao, Y., Chen, S., Lu, W., Gu, W., Liu, P., & Sun, P. (2017). Visible light-induced C3-sulfonamidation of imidazopyridines with sulfamides. *Organic and Biomolecular Chemistry*, 15(38), 8102–8109. <https://doi.org/10.1039/c7ob02029e>
- [116] Gaonkar, S. L., Rai, K. M. L., & Prabhuswamy, B. (2006). Synthesis and antimicrobial studies of a new series of 2-{4-[2-(5-ethylpyridin-2-yl)ethoxy]phenyl}-5-substituted-1,3,4-oxadiazoles. *European Journal of Medicinal Chemistry*, 41(7), 841–846. <https://doi.org/10.1016/j.ejmech.2006.03.002>
- [117] Garcia-Lozano, J., Server-Carrio, J., Escriva, E., Folgado, J.-V., Molla, C., & Lezama, L. (1997). X-ray crystal structure and electronic properties of chlorobis(mepirizole)copper(II) tetrafluoroborate (mepirizole = 4-methoxy-2-(5-methoxy-3-methyl-1H-pyrazol-1-yl)-6-methylpyrimidine). *Polyhedron*, 16(6), 939–944. [https://doi.org/10.1016/S0277-5387\(96\)00346-4](https://doi.org/10.1016/S0277-5387(96)00346-4)

- [118] Gaudreault, J., Varin, F., & Pollack, G. M. (1996). Anticonvulsant Pharmacodynamics and Disposition of Triazolam in Rats These results were presented, in part, at the Seventh Annual Meeting of the American Association of Pharmaceutical Scientists, San Antonio, TX, November 1992. *Journal of Pharmaceutical Sciences*, 85(9), 999–1004. <https://doi.org/10.1021/js9503183>
- [119] Gewald, K. (1965). Heterocyclen aus CH-aciden Nitrilen, VII. 2-Amino-thiophene aus α -Oxo-mercaptanen und methylenaktiven Nitrilen. *Chemische Berichte*, 98(11), 3571–3577. <https://doi.org/10.1002/cber.19650981120>
- [120] Gewald, K., Schinke, E., & Bottcher, H. (1966). Heterocyclen aus CH-aciden Nitrilen, VIII. 2-Amino-thiophene aus methylenaktiven Nitrilen, Carbonylverbindungen und Schwefel. *Chemische Berichte*, 99(1), 94–100. <https://doi.org/10.1002/cber.19660990116>
- [121] Ghaisas, D. S., & Patel, N. B. (2018). Design, Synthesis and Evaluation of Newer Diaryl Ether Analogues Integrated with 1, 3, 4 Oxadiazole Core. *International Journal of Pharmacy and Pharmaceutical Research*, 12(2), 56–67. www.ijppr.humanjournals.com
- [122] Ghanei, S. and L. J. and E. H. and S. M. (2016). Synthesis and Docking Analysis of New Heterocyclic System N1, N4-bis (2-chloroquinolin-3-yl) methylene) benzene-1, 4-diamine as Potential Human AKT1 Inhibitor. *Iranian Journal of Pharmaceutical Research*, 15, 321–327.
- [123] Ghanei, S., Eshghi, H., Lari, J., & Saadatmandzadeh, M. (2015). Synthesis and docking analysis of new 2-chloro-3-((2,2-dimethylhydrazono) methyl)quinoline derivatives as non-nucleoside human HIV-1 reverse transcriptase inhibitors. *Journal of Chemical and Pharmaceutical Research*, 7(2), 428–433. www.jocpr.com
- [124] Girish, Y. R., Kumar, K. S. S., Manasa, H. S., & Shashikanth, S. (2014). ZnO: An Ecofriendly, Green Nano-catalyst for the Synthesis of Pyrazole Derivatives under Aqueous Media. *Journal of the Chinese Chemical Society*, 61(11), 1175–1179. <https://doi.org/10.1002/jccs.201400170>
- [125] Glomb, T., & Swiatek, P. (2021). Antimicrobial activity of 1,3,4-oxadiazole derivatives. *International Journal of Molecular Sciences*, 22(13), 2–23. <https://doi.org/10.3390/ijms22136979>

- [126] Goel, R., Luxami, V., & Paul, K. (2016). Imidazo[1,2-a]pyridines: Promising Drug Candidate for Antitumor Therapy. *Current Topics in Medicinal Chemistry*, 16(30), 3590–3616. <https://doi.org/10.2174/1568026616666160414122644>
- [127] Gomha, S. M. (2009). A facile one-pot synthesis of 6,7,8,9-tetrahydrobenzo[4,5]thieno[2,3-d]-1,2,4-triazolo[4,5-a]pyrimidin-5-ones. *Monatshefte Für Chemie - Chemical Monthly*, 140(2), 213–220. <https://doi.org/10.1007/s00706-008-0060-z>
- [128] Gondek, E., Danel, A., Kwiecień, B., Sanetra, J., & Kityk, A. V. (2009). Optical absorption of bisphenol A based pyrazoloquinoline dimers. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 74(3), 685–690. <https://doi.org/10.1016/j.saa.2009.07.021>
- [129] Gorjizadeh, M., Afshari, M., & Nazari, S. (2013). Microwave-assisted one-step synthesis of 2,5-disubstituted-1,3,4-oxadiazoles using 1,4-bis(triphenylphosphonium)-2-butene peroxodisulfate. *Oriental Journal of Chemistry*, 29(4), 1627–1630. <https://doi.org/10.13005/ojc/290448>
- [130] Greer, N. D. (2003). Voriconazole: The Newest Triazole Antifungal Agent. *Baylor University Medical Center Proceedings*, 16(2), 241–248. <https://doi.org/10.1080/08998280.2003.11927910>
- [131] Griffiths, R. R., Woodson, P. P., & Woodson, P. P. (1988). Reinforcing Properties of Caffeine: Studies in Humans and Laboratory Animals. *Pharmacology Biochemistry & Behavior*, 29.
- [132] Gupta, R., Gupta, A. K., Paul, S., & Kachroo, P. L. (1998). Synthesis and biological activities of some 2-chloro-6/8-substituted-3-(3-alkyl aryl-5,6-dihydro-s-triazolo-[3,4-h] [1,3,4]thiadiazol-6-yl)quinolines. In *Indian Journal of Chemistry* (Vol. 37, pp. 1211–1213).
- [133] Gupta, R. R., Kumar, M., & Gupta, V. (1998). *Heterocyclic Chemistry*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-642-72276-9>
- [134] Hahnor, V. B., Joshi, N. S., Karale, B. K., & Gill, C. H. (2005). Synthesis, characterization of some 1-(2-hydroxy-phenyl)-3-(1-phenyl-3-thiophen-2-yl-1H-pyrazol-4-yl)-propenone, 3-chloro-2-(1-phenyl-3-thiophen-2-yl-1H-pyrazol-4-yl)-chromon-4-one and 2-(1'-phenyl-3'-thiophen-2-yl-3, 4-dihydro-2H,1H'-

- [3,4]bipyrazol-5-yl)-phenol. *Heterocyclic Communications*, 11(2).
<https://doi.org/10.1515/HC.2005.11.2.167>
- [135] Hamed, A. A., Zeid, I. F., El-Ganzory, H. H., & Abdel Aal, M. T. (2008). Synthesis and structure of some thienopyrimidine derivatives. *Monatshefte Fur Chemie*, 139(7), 809–820. <https://doi.org/10.1007/s00706-007-0823-y>
- [136] Hampp, C., Hartzema, A. G., & Kauf, T. L. (2008). Cost-Utility Analysis of Rimonabant in the Treatment of Obesity. *Value in Health*, 11(3), 389–399. <https://doi.org/10.1111/j.1524-4733.2007.00281.x>
- [137] Haque, S. U., Dashwood, M. R., Heetun, M., Shiwen, X., Farooqui, N., Ramesh, B., Welch, H., Savage, F. J., Ogunbiyi, O., Abraham, D. J., & Loizidou, M. (2013). Efficacy of the specific endothelin a receptor antagonist zibotentan (ZD4054) in colorectal cancer: A preclinical study. *Molecular Cancer Therapeutics*, 12(8), 1556–1567. <https://doi.org/10.1158/1535-7163.MCT-12-0975>
- [138] Hartmuth C. Kolb Dr., M. G. F. Prof., K. B. S. P. (2001). Click Chemistry: Diverse Chemical Function from a Few Good Reactions. *German Chemical Society*.
- [139] Hashim, C. S., & Alias, M. F. (2012). Synthesis, spectroscopic study of Pt (IV), Au (III), Rh(III), Co(II) and V (IV) complexes with sodium[5-(p-nitro phenyl)-/ 4-phenyl-1,2,4-triazole-3-dithiocarbamate hydrazide] and cytotoxicity assay on rhabdomyosarcoma cell line of heavy metals. *Baghdad Science Journal*, 9(4), 668–679.
- [140] Hassan, A. F., Abdulwahid, A. T., Al-Luaibi, M. Y., & Al-Jadaan, S. N. (2021). Synthesis, Characterization and Thermal Study of some new Organochalcogenide compounds containing arylamide group. *Egyptian Journal of Chemistry*, 64(9), 5009–5015. <https://doi.org/10.21608/ejchem.2021.49176.3009>
- [141] Hassan, F., & Hameed, A. A. (2014). Synthesis, Characterization and Antioxidant Activity of Some 4-Amino-5-Phenyl-4h-1, 2, 4-Triazole-3-Thiol Derivatives. *International Journal of Applied Science and Technology*, 4(2). <https://www.researchgate.net/publication/323401757>
- [142] Herencia, F., Ferrandiz, M. L., Ubeda, A., Dominguez, Jose N., Charris, J. E., Lobo, G. M., & Alcaraz, M. J. (1998). Synthesis and anti-inflammatory activity of chalcone derivatives. *Bioorganic & Medicinal Chemistry Letters*, 8(10), 1169–1174. [https://doi.org/10.1016/S0960-894X\(98\)00179-6](https://doi.org/10.1016/S0960-894X(98)00179-6)

- [143] Hicks, C., & Gulick, R. M. (2009). Raltegravir: The first HIV Type 1 Integrase Inhibitor. *Clinical Infectious Diseases*, 48(7), 931–939. <https://doi.org/10.1086/597290>
- [144] Hiebel, M. A., Fall, Y., Scherrmann, M. C., & Berteina-Raboin, S. (2014). Straightforward synthesis of various 2,3-diarylimidazo[1,2-a]pyridines in peg400 medium through one-pot condensation and C-H arylation. *European Journal of Organic Chemistry*, 2014(21), 4643–4650. <https://doi.org/10.1002/ejoc.201402079>
- [145] Hierrezuelo, J., Manuel López-Romero, J., Rico, R., Brea, J., Isabel Loza, M., Cai, C., & Algarra, M. (2010). Synthesis of theophylline derivatives and study of their activity as antagonists at adenosine receptors. *Bioorganic and Medicinal Chemistry*, 18(6), 2081–2088. <https://doi.org/10.1016/j.bmc.2010.02.014>
- [146] Holanda, V. N., Silva, W. V. da, Nascimento, P. H. do, Silva, S. R. B., Cabral Filho, P. E., Assis, S. P. de O., Silva, C. A. da, Oliveira, R. N. de, Figueiredo, R. C. B. Q. de, & Lima, V. L. de M. (2020). Antileishmanial activity of 4-phenyl-1-[2-(phthalimido-2-yl)ethyl]-1H-1,2,3-triazole (PT4) derivative on *Leishmania amazonensis* and *Leishmania braziliensis*: In silico ADMET, in vitro activity, docking and molecular dynamic simulations. *Bioorganic Chemistry*, 105, 104437. <https://doi.org/10.1016/j.bioorg.2020.104437>
- [147] Holm, S. C., & Straub, B. F. (2011). Synthesis of n-substituted 1,2,4-triazoles. a review. *Organic Preparations and Procedures International*, 43(4), 319–347. <https://doi.org/10.1080/00304948.2011.593999>
- [148] Hon, K. S., Akolkar, H. N., & Karale, B. K. (2020). Synthesis and characterization of novel 2-(1-benzyl-3-[4-fluorophenyl]-1H-pyrazol-4-yl)-7-fluoro-4H-chromen-4-one derivatives. *Journal of Heterocyclic Chemistry*, 57(4), 1692–1697. <https://doi.org/10.1002/jhet.3894>
- [149] Hsiao, R. H., Tseng, C. C., Xie, J. J., Tsai, S. E., Uramaru, N., Lin, C. Y., Chern, C. Y., & Wong, F. F. (2019). Selective synthesis of functionalized pyrazoles from 5-amino-1H-pyrazole-4-carbaldehydes with sodium nitrite: 5-Amino-4-nitrosopyrazoles and pyrazole-4-carbaldehydes. *Tetrahedron*, 75(33), 4561–4569. <https://doi.org/10.1016/j.tet.2019.06.048>
- [150] Huang, J., Wu, D., Bai, X., Cai, P., & Zhu, W. G. (2021). Catalyst-free, visible-light-induced direct radical cross-coupling perfluoroalkylation of the imidazo[1,2-

- a]pyridines with perfluoroalkyl iodides. *New Journal of Chemistry*, 45(11), 4925–4929. <https://doi.org/10.1039/d1nj00651g>
- [151] Huang, Q., Dong, H., Li, B., Hu, W., & Wang, Y. (2020). Rhodium catalyzed direct C3-ethoxycarbonylmethylation of imidazo[1,2-a]pyridines with ethyl diazoacetate. *Tetrahedron*, 76(10). <https://doi.org/10.1016/j.tet.2020.130998>
- [152] Huisgen, R. (1963). 1,3-Dipolar Cycloadditions. Past and Future. In *Angewandte Chemie International Edition in English* (Vol. 2, Issue 10). <https://doi.org/10.1002/anie.196305651>
- [153] Husain, A., Ahuja, P., & Sarafroz. (2009). Synthesis and biological evaluation of β -aroylpropionic acid based 1,3,4-oxadiazoles. *Indian Journal of Pharmaceutical Sciences*, 71(1), 62. <https://doi.org/10.4103/0250-474X.51963>
- [154] Husain, A., & Ajmal, M. (2009). Synthesis of novel 1,3,4-oxadiazole derivatives and their biological properties. *Acta Pharmaceutica*, 59(2), 223–233. <https://doi.org/10.2478/v10007-009-0011-1>
- [155] Hussain, A., Sharba, K., Al-Bayati, R. H., Aouad, M., & Rezki, N. (2005). molecules Synthesis of Oxadiazoles, Thiadiazoles and Triazoles Derived from Benzo[b]thiophene. *Molecules*, 10, 1161–1168. <http://www.mdpi.org>
- [156] Hutton, S. M., Mackay, S. P., & Meth-Cohn, O. (2000). Synthesis of Dibenzo[c,h][1,6]naphthyridine, [2]Benzopyrano[4,3-c]quinoline and Benzo[i]phenanthridine Analogues of the Quaternary Benzo[c]phenanthridines. *Synthesis*, 2000(08), 1121–1124. <https://doi.org/10.1055/s-2000-6325>
- [157] Ibrahim, N. M., A Yosef, H. A., & Mahran, M. H. (2010). *The Behavior of 2-Chloroquinoline-3-carboxaldehyde towards Certain Primary Amines and Activated Methylenes* (Vol. 53, Issue 5).
- [158] Ingale, S., & Dawson, P. E. (2011). On Resin Side-Chain Cyclization of Complex Peptides Using CuAAC. *Organic Letters*, 13(11), 2822–2825. <https://doi.org/10.1021/ol200775h>
- [159] Isloor, A. M., Kalluraya, B., & Shetty, P. (2009). Regioselective reaction: Synthesis, characterization and pharmacological studies of some new Mannich bases derived from 1,2,4-triazoles. *European Journal of Medicinal Chemistry*, 44(9), 3784–3787. <https://doi.org/10.1016/j.ejmech.2009.04.038>

- [160] Ismail, M. A., Arafa, R. K., Wenzler, T., Brun, R., Tanious, F. A., Wilson, W. D., & Boykin, D. W. (2008). Synthesis and antiprotozoal activity of novel bis-benzamidino imidazo[1,2-a]pyridines and 5,6,7,8-tetrahydro-imidazo[1,2-a]pyridines. *Bioorganic and Medicinal Chemistry*, 16(2), 683–691. <https://doi.org/10.1016/j.bmc.2007.10.042>
- [161] Ismail, M. A. H., Lehmann, J., Abou El Ella, D. A., Albohy, A., & Abouzid, K. A. M. (2009). Lonazolac analogues: molecular modeling, synthesis, and in vivo anti-inflammatory activity. *Medicinal Chemistry Research*, 18(9), 725–744. <https://doi.org/10.1007/s00044-009-9163-2>
- [162] Jacobs, M. R., & Appelbaum, P. C. (2006). Nadifloxacin: a quinolone for topical treatment of skin infections and potential for systemic use of its active isomer, WCK 771. In *Expert Opinion on Pharmacotherapy* (Vol. 7, Issue 14). <https://doi.org/10.1517/14656566.7.14.1957>
- [163] Jairaj Kishanrao, D., Madhuri Kumar, P., Sushil Ramakant, M., Kalimoddin Inayatsab, omin, Abhay Shivaji, B., Research Center, P. G., & Madhavarao Patil Mahavidhyalaya, S. (2018). Design and synthesis series of novel quinolinyl methanol's derivatives. *World Journal of Pharmacy and Pharmaceutical Sciences*, 2(5), 3373–3381. www.wjpps.com
- [164] Jalal, M. A. F., & Collin, H. A. (1976). Estimation of caffeine, theophylline and theobromine in plant material. *New Phytologist*, 76(2), 277–281. <https://doi.org/10.1111/j.1469-8137.1976.tb01461.x>
- [165] Jassim, I. K., Kh Jassim, W., Abd ALsatar, S., & HMohammed, A. (2012). Synthesis and characterization of some new of thiazolidine ,1,2,4-triazole ,1,3,4-thiadiazole , semicarbazide , oxazoline and a study of their biological activity. . *Kerbala Journal of Pharmaceutical Sciences Number*, 3(3), 213–222.
- [166] Javahershenas, R., & Nikzat, S. (2024). Recent developments using malononitrile in ultrasound-assisted multicomponent synthesis of heterocycles. *Ultrasonics Sonochemistry*, 102, 106741. <https://doi.org/10.1016/j.ultsonch.2023.106741>
- [167] Jaware, J., & Borhade, S. (2013). Synthesis of novel N-{[2-(morpholin-4-yl)-quinoline-3-yl]methyl}2-[(4-aminopentyl)(ethyl)amino]ethanol derivatives. *International Journal of Pharmacy and Pharmaceutical Reseach*, 23(2), 67–71.
- [168] Jia, Z., Wei, S., & Zhu, Q. (2010). Monoamine Oxidase Inhibitors: Benzyldiene-prop-2-ynyl-amines Analogues Fig. 1. Proposed Mechanism of Covalent Inactivation

- of MAO by Propar-gylamine Inhibitors. In *Biol. Pharm. Bull* (Vol. 33, Issue 4, pp. 725–728).
- [169] Johnson, G. F., Tu, J.-I., Bartlett, M. L. S., & Graves, D. J. (1970). Physical-Chemical Studies on the Pyridoxal Phosphate Binding Site in Sodium Borohydride-reduced and Native Phosphorylase. *Journal of Biological Chemistry*, 245(21), 5560–5568. [https://doi.org/10.1016/S0021-9258\(18\)62692-4](https://doi.org/10.1016/S0021-9258(18)62692-4)
- [170] Jones, G. O., & Houk, K. N. (2008). Predictions of Substituent Effects in Thermal Azide 1,3-Dipolar Cycloadditions: Implications for Dynamic Combinatorial (Reversible) and Click (Irreversible) Chemistry. *The Journal of Organic Chemistry*, 73(4), 1333–1342. <https://doi.org/10.1021/jo702295d>
- [171] Joshi, D., Narigara, R., Jani, G., & Parikh, K. (2019). Synthesis and Evaluation of Novel Fluorobenzimidazole Derivatives as Antibacterial and Antifungal Agents. *Asian Journal of Organic & Medicinal Chemistry*, 4(4), 209–215. <https://doi.org/10.14233/ajomc.2019.ajomc-p200>
- [172] Joshi, S. & J. A. & V. H. M. & V. V. P. & G. G. S. . (2010). Synthesis of some new quinolinylpyrazoles as potential antibacterial and antitubercular agents. *Indian Journal of Heterocyclic Chemistry*, 19, 221–224.
- [173] Joshi, S. D., More, U. A., Parkale, D., Aminabhavi, T. M., Gadad, A. K., Nadagouda, M. N., & Jawarkar, R. (2015). Design, synthesis of quinolinyl Schiff bases and azetidinones as enoyl ACP-reductase inhibitors. *Medicinal Chemistry Research*, 24(11), 3892–3911. <https://doi.org/10.1007/s00044-015-1432-7>
- [174] Julino, M., & Stevens, M. F. G. (1998). Antitumour polycyclic acridines. Part 5.1 Synthesis of 7H-pyrido[4,3,2-kl]acridines with exploitable functionality in the pyridine ring. *Journal of the Chemical Society, Perkin Transactions 1*, 10, 1677–1684. <https://doi.org/10.1039/a800575c>
- [175] Kamble, R., & Sudha, B. (2006). Synthesis and pharmacological screening of 5-methyl-3-[p-(6'-aryl-2'-thioxo-1',2',5',6'-tetrahydro-pyrimidin-4'-yl)-phenyl]-3H -2-oxo-4d-1,3,4-oxadiazoles. *Indian Journal of Pharmaceutical Sciences*, 68(2), 249. <https://doi.org/10.4103/0250-474X.25729>
- [176] Kaneriya, D. J., Vadodaria, M. S., Patel, A. U., & Ladva, K. D. (2016). Chemical Science Review and Letters Synthesis of Some Chalcones and Pyrazoline bearing

Chloroquinoline Scaffold as Biologically Active Agents. *Chem Sci Rev Lett*, 5(17), 1–4.

- [177] Kapupara, V. H., Kalavadiya, P. L., Gojiya, D. G., Parmar, N. D., & Joshi, H. S. (2019). Greener, facile and ultrasound assisted synthesis of 1,2,3-triazole via CuAAC of 2-(azidomethyl)-1H-benzo[d]imidazole and alkynes and their evaluation of antimicrobial activity. *International Scientific Journal*, 119, 139–167. www.worldscientificnews.com
- [178] Karale, U. B., Kalari, S., Shivakumar, J., Makane, V. B., Babar, D. A., Thakare, R. P., Babu, B. N., Chopra, S., & Rode, H. B. (2016). Ligand-free Pd-catalysed decarboxylative arylation of imidazo[1,2-a] pyridine-3-carboxylic acids with aryl bromides. *RSC Advances*, 6(69), 65095–65104. <https://doi.org/10.1039/c6ra12166g>
- [179] Karthikeyan, M. S., & Holla, B. S. (2008). Synthesis, antiinflammatory and antimicrobial activities of some 2,4-dichloro-5-fluorophenyl substituted arylidenetriazolothiazolidinones. *Monatshefte Für Chemie - Chemical Monthly*, 139(6), 691–696. <https://doi.org/10.1007/s00706-007-0794-z>
- [180] Kaur, K., Jain, M., Reddy, R. P., & Jain, R. (2010a). Quinolines and structurally related heterocycles as antimalarials. *European Journal of Medicinal Chemistry*, 45(8), 3245–3264. <https://doi.org/10.1016/j.ejmech.2010.04.011>
- [181] Kaur, K., Jain, M., Reddy, R. P., & Jain, R. (2010b). Quinolines and structurally related heterocycles as antimalarials. *European Journal of Medicinal Chemistry*, 45(8), 3245–3264. <https://doi.org/10.1016/j.ejmech.2010.04.011>
- [182] Kaur, R., Dwivedi, A. R., Kumar, B., & Kumar, V. (2016). Recent Developments on 1,2,4-Triazole Nucleus in Anticancer Compounds: A Review. *Anti-Cancer Agents in Medicinal Chemistry*, 16(4), 465–489.
- [183] Kaushik, C. P., Luxmi, R., Singh, D., & Kumar, A. (2017). Synthesis and antimicrobial evaluation of ester-linked 1,4-disubstituted 1,2,3-triazoles with a furyl/thienyl moiety. *Molecular Diversity*, 21(1), 137–145. <https://doi.org/10.1007/s11030-016-9710-y>
- [184] Ke, S., Li, Z., & Qian, X. (2008). 1,3,4-Oxadiazole-3(2H)-carboxamide derivatives as potential novel class of monoamine oxidase (MAO) inhibitors: Synthesis, evaluation, and role of urea moiety. *Bioorganic & Medicinal Chemistry*, 16(16), 7565–7572. <https://doi.org/10.1016/j.bmc.2008.07.026>

- [185] Keri, R. S., & Patil, S. A. (2014). Quinoline: A promising antitubercular target. *Biomedicine & Pharmacotherapy*, 68(8), 1161–1175. <https://doi.org/10.1016/j.biopha.2014.10.007>
- [186] Khalil, S. M. E. (2003). Synthesis, Spectroscopic and Magnetic Studies on Metal Complexes of 5-Methyl-3-(2-Hydroxyphenyl)Pyrazole. *Journal of Coordination Chemistry*, 56(12), 1013–1024. <https://doi.org/10.1080/0095897031000135289>
- [187] Khan, M. F., Alam, M. M., Verma, G., Akhtar, W., Akhter, M., & Shaquiquzzaman, M. (2016). The therapeutic voyage of pyrazole and its analogs: A review. *European Journal of Medicinal Chemistry*, 120, 170–201. <https://doi.org/10.1016/j.ejmech.2016.04.077>
- [188] Khidre, R. E., Abu-Hashem, A. A., & El-Shazly, M. (2011). Synthesis and antimicrobial activity of some 1- substituted amino-4,6-dimethyl-2-oxo-pyridine-3-carbonitrile derivatives. *European Journal of Medicinal Chemistry*, 46(10), 5057–5064. <https://doi.org/10.1016/j.ejmech.2011.08.018>
- [189] Khillare, L., Bhosle, M., Bhalerao, M., Kharat, K., & Mane, R. (2017). Synthesis of New Thiazolyl Coupled Pyrazoles Bearing 2,4-Thiazolidinedionyl Pharmacophore and Their Anti-Inflammatory and Antibacterial Evaluation. *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry*, 16(1), 46–57. <https://doi.org/10.2174/1871523016666170616120346>
- [190] Kidwai, M., Saxena, S., Rahman Khan, M. K., & Thukral, S. S. (2005). Aqua mediated synthesis of substituted 2-amino-4H-chromenes and in vitro study as antibacterial agents. *Bioorganic & Medicinal Chemistry Letters*, 15(19), 4295–4298. <https://doi.org/10.1016/j.bmcl.2005.06.041>
- [191] Kingsbury, W. D., Boehm, J. C., Jakas, D. R., Holden, K. G., Hecht, S. M., Gallagher, G., Caranfa, M. J., McCabe, F. L., Faucette, L. F., Johnson, R. K., & Hertzberg, R. P. (1991). Synthesis of water-soluble (aminoalkyl)camptothecin analogs: inhibition of topoisomerase I and antitumor activity. *Journal of Medicinal Chemistry*, 34(1), 98–107. <https://doi.org/10.1021/jm00105a017>
- [192] Kiyani, H., Albooyeh, F., & Fallahnezhad, S. (2015). Synthesis of new pyrazolyl-1,3-diazabicyclo[3.1.0]hexe-3-ene derivatives. *Journal of Molecular Structure*, 1091, 163–169. <https://doi.org/10.1016/j.molstruc.2015.02.069>
- [193] *knorr1883*. (n.d.).

- [194] Kocyigit, U. M., Taslimi, P., Tuzun, B., Yakan, H., Muglu, H., & Guzel, E. (2022). 1,2,3-Triazole substituted phthalocyanine metal complexes as potential inhibitors for anticholinesterase and antidiabetic enzymes with molecular docking studies. *Journal of Biomolecular Structure and Dynamics*, 40(10), 4429–4439. <https://doi.org/10.1080/07391102.2020.1857842>
- [195] Kommidi, H., Guo, H., Nurili, F., Vedvyas, Y., Jin, M. M., McClure, T. D., Ehdaie, B., Sayman, H. B., Akin, O., Aras, O., & Ting, R. (2018). 18 F-Positron Emitting/Trimethine Cyanine-Fluorescent Contrast for Image-Guided Prostate Cancer Management. *Journal of Medicinal Chemistry*, 61(9), 4256–4262. <https://doi.org/10.1021/acs.jmedchem.8b00240>
- [196] Koparr, M., Cetin, A., & Cansız, A. (2005). 5-Furan-2yl[1,3,4]oxadiazole-2-thiol, 5-Furan-2yl-4H [1,2,4] triazole-3-thiol and Their Thiol-Thione Tautomerism. *Molecules*, 10(2), 475–480. <https://doi.org/10.3390/10020475>
- [197] Krishna Prasad, C., & Machiraju, P. V. S. (2017). Synthesis and characterization of some novel aryl and heteroaryl chalcone derivatives of 3-(3,4,5-trimethoxyphenyl)-1-phenyl-1H-pyrazole-4-carbaldehyde for assessing their potentials as anticancer agents. *Russian Journal of General Chemistry*, 87(9), 2056–2066. <https://doi.org/10.1134/S1070363217090225>
- [198] Krishnamoorthy, R., & Anaikutti, P. (2023). Iodine catalyzed synthesis of imidazo[1,2-a]pyrazine and imidazo[1,2-a]pyridine derivatives and their anticancer activity. *Royal Society of Advances*, 13(51), 36439–36454. <https://doi.org/10.1039/D3RA07842F>
- [199] Kulabas, N., Tatar, E., Bingol ozakpinar, ozlem, ozsavcı, D., Pannecouque, C., De Clercq, E., & Kucukguzel, ilkay. (2016). Synthesis and antiproliferative evaluation of novel 2-(4H-1,2,4-triazole-3-ylthio)acetamide derivatives as inducers of apoptosis in cancer cells. *European Journal of Medicinal Chemistry*, 121, 58–70. <https://doi.org/10.1016/j.ejmech.2016.05.017>
- [200] Kumar, R., Rawat, D., Semwal, R., Badhani, G., & Adimurthy, S. (2021). Hypervalent iodine mediated synthesis of imidazo[1,2-: A] pyridine ethers: Consecutive methylene linkage and insertion of ethylene glycol. *New Journal of Chemistry*, 45(17), 7491–7495. <https://doi.org/10.1039/d1nj00657f>

- [201] Kumar Shukla, P., Soni, N., Verma, A., & Kumar Jha, A. (2014). Synthesis, characterization and in vitro biological evaluation of a series of 1,2,4-triazoles derivatives & triazole based schiff bases. *Scholars Research Library Der Pharma Chemica*, 6(3), 153–160. <http://derpharmachemica.com/archive.html>
- [202] Kumar Singh, A., & Raj Kandel, K. (2012). Synthesis of Triazole derivative: [4-(benzylideneamino)-5-phenyl-4H-1,2,4-triazole-3-thiol]. *Journal of Nepal Chemical Society*, 30, 174–177.
- [203] Kumari, M. A., Rao, C. V., Triloknadh, S., Harikrishna, N., Venkataramaiah, C., Rajendra, W., Trinath, D., & Suneetha, Y. (2018). Synthesis, docking and ADME prediction of novel 1,2,3-triazole-tethered coumarin derivatives as potential neuroprotective agents. *Research on Chemical Intermediates*, 44(3), 1989–2008. <https://doi.org/10.1007/s11164-017-3210-2>
- [204] Kurteva, V. B., Lubenov, L. A., Nedeltcheva, D. V, Nikolova, R. P., & Shivachev, B. L. (2012). *Fast and efficient direct conversion of 2-aminopyridine into 2,3-disubstituted imidazo[1,2-a]pyridines*.
- [205] Kusy, D., Marchwicka, A., Małolepsza, J., Justyna, K., Gendaszewska-Darmach, E., & Błażewska, K. M. (2021). Synthesis of the 6-Substituted Imidazo[1,2-a]Pyridine-3-yl-2- Phosphonopropionic Acids as Potential Inhibitors of Rab Geranylgeranyl Transferase. *Frontiers in Chemistry*, 8. <https://doi.org/10.3389/fchem.2020.596162>
- [206] Kwong, H. C., Chidan Kumar, C. S., Mah, S. H., Mah, Y. L., Chia, T. S., Quah, C. K., Lim, G. K., & Chandraju, S. (2019). Crystal Correlation Of Heterocyclic Imidazo[1,2-a]pyridine Analogues and Their Anticholinesterase Potential Evaluation. *Scientific Reports*, 9(1), 926. <https://doi.org/10.1038/s41598-018-37486-7>
- [207] Lamani, D. S., Venugopala Reddy, K. R., Bhojya Naik, H. S., Savyasachi, A., & Naik, H. R. (2008). Synthesis and DNA binding studies of novel heterocyclic substituted quinoline schiff bases: A potent antimicrobial agent. *Nucleosides, Nucleotides and Nucleic Acids*, 27(10–11), 1197–1210. <https://doi.org/10.1080/15257770802400081>
- [208] Lee, L. V., Mitchell, M. L., Huang, S.-J., Fokin, V. V., Sharpless, K. B., & Wong, C.-H. (2003). A Potent and Highly Selective Inhibitor of Human α -1,3-Fucosyltransferase via Click Chemistry. *Journal of the American Chemical Society*, 125(32), 9588–9589. <https://doi.org/10.1021/ja0302836>

- [209] Li, C. R., Liu, Z. C., Wang, B. D., Li, T. R., & Yang, Z. Y. (2015). Synthesis and fluorescence properties of Sm-2-oxo-quinoline-3-carbaldehyde-isonicotinyl hydrazone decorated with 1,10-phenanthroline. *Synthetic Metals*, 209, 273–278. <https://doi.org/10.1016/j.synthmet.2015.08.009>
- [210] Li, J., Li, Z., Li, T., Lin, L., Zhang, Y., Guo, L., Xu, Y., Zhao, W., & Wang, P. (2012). Identification of a specific inhibitor of nOGA — a caspase-3 cleaved O-GlcNAcase variant during apoptosis. *Biochemistry (Moscow)*, 77(2), 194–200. <https://doi.org/10.1134/S0006297912020113>
- [211] Li, M., & Zhao, B.-X. (2014). Progress of the synthesis of condensed pyrazole derivatives. *European Journal of Medicinal Chemistry*, 85, 311–340. <https://doi.org/10.1016/j.ejmech.2014.07.102>
- [212] Li, T., Wen, G., Li, J., Zhang, W., & Wu, S. (2019). A useful synthesis of 2-acylamino-1,3,4-oxadiazoles from acylthiosemicarbazides using potassium iodate and the discovery of new antibacterial compounds. *Molecules*, 24(8). <https://doi.org/10.3390/molecules24081490>
- [213] Li, X., Wang, S., Zang, J., Liu, M., Jiang, G., & Ji, F. (2020). Copper- And DMF-mediated switchable oxidative C-H cyanation and formylation of imidazo[1,2-: A] pyridines using ammonium iodide. *Organic and Biomolecular Chemistry*, 18(44), 9100–9108. <https://doi.org/10.1039/d0ob01838d>
- [214] Liu, Z. C., Wang, B. D., Li, B., Wang, Q., Yang, Z. Y., Li, T. R., & Li, Y. (2010). Crystal structures, DNA-binding and cytotoxic activities studies of Cu(II) complexes with 2-oxo-quinoline-3-carbaldehyde Schiff-bases. *European Journal of Medicinal Chemistry*, 45(11), 5353–5361. <https://doi.org/10.1016/j.ejmech.2010.08.060>
- [215] Lo, Y.-C., Tsou, H.-H., Lin, R.-J., Wu, D.-C., Wu, B.-N., Lin, Y.-T., & Chen, I.-J. (2005). Endothelium-dependent and -independent vasorelaxation by a theophylline derivative MCPT: Roles of cyclic nucleotides, potassium channel opening and phosphodiesterase inhibition. *Life Sciences*, 76(8), 931–944. <https://doi.org/10.1016/j.lfs.2004.10.009>
- [216] Lu, L., Wu, J., & Yang, C. (2008). Preparation of 1H-1,2,3-Triazoles by Cuprous Ion Mediated Cycloaddition of Terminal Alkyne and Sodium Azide. *Journal of the Chinese Chemical Society*, 55(2), 414–417. <https://doi.org/10.1002/jccs.200800061>

- [217] Luan, N., Liu, Z., Han, S., Shen, L., Li, J., Zou, D., Wu, Y., & Wu, Y. (2020). PhI(OAc)₂-mediated oxidative C[α]H sulfoximation of imidazopyridines under mild conditions. *Tetrahedron Letters*, 61(1). <https://doi.org/10.1016/j.tetlet.2019.151362>
- [218] Lyon, M. A., Lawrence, S., Williams, D. J., & Jackson, Y. A. (1999). Synthesis and structure verification of an analogue of kuanoniamine A. *Journal of the Chemical Society, Perkin Transactions 1*, 4, 437–442. <https://doi.org/10.1039/a809203f>
- [219] Ma, L., Wang, X., Yu, W., & Han, B. (2011). TBAI-catalyzed oxidative coupling of aminopyridines with β -keto esters and 1,3-diones - Synthesis of imidazo[1,2-a]pyridines. *Chemical Communications*, 47(40), 11333–11335. <https://doi.org/10.1039/c1cc13568f>
- [220] Ma, L. Y., Pang, L. P., Wang, B., Zhang, M., Hu, B., Xue, D. Q., Shao, K. P., Zhang, B. Le, Liu, Y., Zhang, E., & Liu, H. M. (2014). Design and synthesis of novel 1,2,3-triazole-pyrimidine hybrids as potential anticancer agents. *European Journal of Medicinal Chemistry*, 86, 368–380. <https://doi.org/10.1016/j.ejmech.2014.08.010>
- [221] Maghraby, M. T. E., Mazyad Almutairi, T., Bräse, S., Salem, O. I. A., Youssif, B. G. M., & Sheha, M. M. (2023). New 1,2,3-Triazole/1,2,4-triazole Hybrids as Aromatase Inhibitors: Design, Synthesis, and Apoptotic Antiproliferative Activity. *Molecules*, 28(20). <https://doi.org/10.3390/molecules28207092>
- [222] Maguire, M. P., Sheets, K. R., McVety, K., Spada, A. P., & Zilberstein, A. (1994). A New Series of PDGF Receptor Tyrosine Kinase Inhibitors: 3-Substituted Quinoline Derivatives. *Journal of Medicinal Chemistry*, 37(14), 2129–2137. <https://doi.org/10.1021/jm00040a003>
- [223] Mailavaram Raghu Prasad, a A. R. R. R. P. S. R. K. S. R. (2001). Microwave-Assisted Synthesis of Novel 5-Substituted-2,3-dihydroimidazo[1,2-c]thieno[3,2-e]pyrimidines. *Thieme Stuttgart*, 14, 2119–2123.
- [224] Malcolm Sainsbury. (1992). *Rodd's Chemistry of carbon compounds* (sainsbury malcoms, Ed.; 2nd ed., Vol. 1).
- [225] Malladi, S., Isloor, A. M., Isloor, S., Akhila, D. S., & Fun, H.-K. (2013). Synthesis, characterization and antibacterial activity of some new pyrazole based Schiff bases. *Arabian Journal of Chemistry*, 6(3), 335–340. <https://doi.org/10.1016/j.arabjc.2011.10.009>

- [226] Mallandur, B. K., Rangaiah, G., & Harohally, N. V. (2017). Synthesis and antimicrobial activity of Schiff bases derived from 2-chloro quinoline-3-carbaldehyde and its derivatives incorporating 7-methyl-2-propyl-3H-benzimidazole-5-carboxylic acid hydrazide. *Synthetic Communications*, 47(11), 1065–1070. <https://doi.org/10.1080/00397911.2017.1309668>
- [227] Manap, S., Gursay-Kol, O., Alkan, M., & Yuksek, H. (2020). Synthesis, in vitro antioxidant and antimicrobial activities of some novel 3-substituted-4-(3-methoxy-4-isobutyryloxybenzylideneamino)-4,5-dihydro-1H-1,2,4-triazol-5-one derivatives. *Indian Journal of Chemistry*, 59B, 271–282.
- [228] Mancinelli, A., Guiso, G., Garattini, S., Urso, R., & Caccia, S. (1985). Kinetic and pharmacological studies on estazolam in mice and man. *Xenobiotica*, 15(3), 257–265. <https://doi.org/10.3109/00498258509045357>
- [229] Mantzanidou, M., Pontiki, E., & Hadjipavlou-Litina, D. (2021). Pyrazoles and Pyrazolines as Anti-Inflammatory Agents. *Molecules*, 26(11), 3439. <https://doi.org/10.3390/molecules26113439>
- [230] Marouf Al-Azzawi, A., & Khalaf Hammud, K. (2013). Synthesis and Characterization of Some New 1,3,4-Oxadiazole and 1,2,4-Triazole Derivatives Based on 3,4,5,6 Tetrachlorophthalimide. *Iraqi Journal of Science*, 54(4), 782–788.
- [231] Martin, P. J., & Bruce, D. W. (2007). Hydrogen-bonded oxadiazole mesogens. *Liquid Crystals*, 34(6), 767–774. <https://doi.org/10.1080/02678290701343273>
- [232] Matiichuk, V. S., Potopnyk, M. A., & Obushak, N. D. (2008). Molecular design of pyrazolo[3,4-d]pyridazines. *Russian Journal of Organic Chemistry*, 44(9), 1352–1361. <https://doi.org/10.1134/S1070428008090182>
- [233] Matsumura, M., Takahashi, T., Yamauchi, H., Sakuma, S., Hayashi, Y., Hyodo, T., Obata, T., Yamaguchi, K., Fujiwara, Y., & Yasuike, S. (2020). Synthesis and anticancer activity of bis(2-arylimidazo[1,2-a]pyridin-3-yl) selenides and diselenides: The copper-catalyzed tandem C-H selenation of 2-arylimidazo[1,2-a]pyridine with selenium. *Beilstein Journal of Organic Chemistry*, 16, 1075–1083. <https://doi.org/10.3762/bjoc.16.94>
- [234] Mazurek, A. P., Bojarski, J., Marcinkowski, K., Furmanowa, M., Gutkowska, B., Kaliszan, R., Pachecka, J., Pawlaczyk, J., Pluta, J., Wieniawski, W., Komarek, P., Ostrowski-Meissner, H., R^der, E., Skolnick, P., & Vincze, Z. (2014). Synthesis and

- anti-mycobacterial activity of new 4-thiazolidinone and 1,3,4-oxadiazole derivatives of isoniazid. *Drug Research ACTA POLONIAE PHARMACEUTICA*, 71, 763–770. www.actapoloniaepharmaceutica.pl
- [235] Medapi, B., Renuka, J., Saxena, S., Sridevi, J. P., Medishetti, R., Kulkarni, P., Yogeeswari, P., & Sriram, D. (2015). Design and synthesis of novel quinoline–aminopiperidine hybrid analogues as Mycobacterium tuberculosis DNA gyraseB inhibitors. *Bioorganic & Medicinal Chemistry*, 23(9), 2062–2078. <https://doi.org/10.1016/j.bmc.2015.03.004>
- [236] Meher, D. K., Nayak, A. K., Singh, A. K., & Pathak, M. (2020). Synthesis, Evaluation and Pharmacological action of Oxadiazole derivatives. *Journal of Pharmacognosy and Phytochemistry*, 9(6), 1307–1315. <https://doi.org/10.22271/phyto.2020.v9.i6s.13132>
- [237] Meth-Cohn, O. (1993). The synthesis of pyridines, quinolines and other related systems by the vilsmeier and the reverse vilsmeier method. *Heterocycles*, 35(1), 539–557.
- [238] Meth-Cohn, O., Narine, B., & Tarnowski, B. (1981). A versatile new synthesis of quinolines and related fused pyridines. Part 5. The synthesis of 2-chloroquinoline-3-carbaldehydes. *Journal of the Chemical Society, Perkin Transactions 1*, 1520–1530. <https://doi.org/10.1039/P19810001520>
- [239] Michael, J. P. (2004). Quinoline, quinazoline and acridone alkaloids. *Natural Product Reports*, 21(5), 650. <https://doi.org/10.1039/b310691h>
- [240] Miniyaar, P. B., Barmade, M. A., & Mahajan, A. A. (2015). Synthesis and biological evaluation of 1-(5-(2-chloroquinolin-3-yl)-3-phenyl-1H-pyrazol-1-yl)ethanone derivatives as potential antimicrobial agents. *Journal of Saudi Chemical Society*, 19(6), 655–660. <https://doi.org/10.1016/j.jscs.2013.12.004>
- [241] Miri, R., Razzaghi-asl, N., & Mohammadi, M. K. (2013). QM study and conformational analysis of an isatin Schiff base as a potential cytotoxic agent. *Journal of Molecular Modeling*, 19(2), 727–735. <https://doi.org/10.1007/s00894-012-1586-x>
- [242] Mistry, B. M., & Jauhari, S. (2013). Synthesis and in vitro antimicrobial and anti-tubercular evaluation of some quinoline-based azitidinone and thiazolidinone analogues. *Medicinal Chemistry Research*, 22(2), 635–646. <https://doi.org/10.1007/s00044-012-0060-8>

- [243] Mobinikhaledi, A., Foroughifar, N., Khanpour, M., & Ebrahimi, S. (2010). Synthesis of some novel Schiff bases containing 1,2,4-triazole ring. *European Journal of Chemistry*, 1(1), 33–36. <https://doi.org/10.5155/eurjchem.1.1.33-36.5>
- [244] Mohammadi-Khanaposhtani, M., Nori, M., Valizadeh, Y., Javanshir, S., Dastyafteh, N., Moaazam, A., Hosseini, S., Larijani, B., Adibi, H., Biglar, M., Hamedifar, H., Mahdavi, M., Kamci, H., Karakus, A., & Taslimi, P. (2022). New 4-phenylpiperazine-carbodithioate- N-phenylacetamide hybrids: Synthesis, in vitro and in silico evaluations against cholinesterase and α -glucosidase enzymes. *Archiv Der Pharmazie*, 355(5). <https://doi.org/10.1002/ardp.202100313>
- [245] Mohammed, H. (2021). Synthesis, Identification, and Biological Study for Some Complexes of Azo Dye Having Theophylline. *The Scientific World Journal*, 2021, 1–9. <https://doi.org/10.1155/2021/9943763>
- [246] Molina, P., Arques, A., Vinader, M. V., Becher, J., & Brondum, K. (1988). Fused pyrimidines by a tandem aza-Wittig/electrocyclic ring closure strategy: synthesis of pyrazolo[3,4-d]pyrimidine, [1,2,3]triazolo[4,5-d]pyrimidine, and thiazolo[4,5-d]pyrimidine derivatives. *The Journal of Organic Chemistry*, 53(20), 4654–4663. <https://doi.org/10.1021/jo00255a003>
- [247] Moraski, G. C., Markley, L. D., Hipskind, P. A., Boshoff, H., Cho, S., Franzblau, S. G., & Miller, M. J. (2011). Advent of imidazo[1,2-a]pyridine-3-carboxamides with potent multi- and extended drug resistant antituberculosis activity. *ACS Medicinal Chemistry Letters*, 2(6), 466–470. <https://doi.org/10.1021/ml200036r>
- [248] Mosselhi, M. A. N., Darwish, E. S., & Peseke, K. (2008). Synthesis and antimicrobial activity of imidazo- and pyrimido[2,1-f]-theophyllines. *Monatshefte Für Chemie*, 139(7), 825–834. <https://doi.org/10.1007/s00706-008-0854-z>
- [249] Moulin, A., Bibian, M., Blayo, A. L., El Habnoui, S., Martinez, J., & Fehrentz, J. A. (2010). Synthesis of 3,4,5-trisubstituted-1,2,4-triazoles. *Chemical Reviews*, 110(4), 1809–1827. <https://doi.org/10.1021/cr900107r>
- [250] Mounika, K. , P. A. , & G. C. (2010). Synthesis characterization and biological activity of a Schiff base derived from 3-ethoxy salicylaldehyde and 2-amino benzoic acid and its transition metal complexes. *Journal of Scientific Research*, 2(3), 513.
- [251] Moureu, C. and D. R. (1901). Over some Acetylenketone and over a new method to the synthesis of beta-Diketones. *Bull. Soc. Chim. Fr*, 25(1), 302–3013.

- [252] Mphahlele, M., & Adeloje, A. (2013). 4,6,8-Triarylquinoline-3-carbaldehyde Derivatives: Synthesis and Photophysical Properties. *Molecules*, 18(12), 15769–15787. <https://doi.org/10.3390/molecules181215769>
- [253] Muddam, B., Venkanna, P., Venkateswarlu, M., Satish Kumar, M., & Rajanna, K. (2018). Symmetrical Trichlorotriazine Derivatives as Efficient Reagents for One-Pot Synthesis of 3-Acetyl-2-chloroquinolines from Acetanilides under Vilsmeier–Haack Conditions. *Synlett*, 29(01), 85–88. <https://doi.org/10.1055/s-0036-1589099>
- [254] Mustafa, A. M., Sayyid, F. F., Betti, N., shaker, L. M., Hanoon, M. M., Alamiery, A. A., Kadhum, A. A. H., & Takriff, M. S. (2022). Inhibition of mild steel corrosion in hydrochloric acid environment by 1-amino-2-mercapto-5-(4-(pyrrol-1-yl)phenyl)-1,3,4-triazole. *South African Journal of Chemical Engineering*, 39, 42–51. <https://doi.org/10.1016/j.sajce.2021.11.009>
- [255] N. Sampathkumar, N. V. K. and S. P. R. (2004). A simple synthesis of dibenzo[b,g][1,8]naphthyridines. *Synthetic Communications*, 34(11), 9–16. <https://doi.org/10.1081/scc-120037926>
- [256] Nabeshima, T., Ina, M., & Kameyama, T. (1977). Effect of mono-amine-related drugs on inhibitory action of difenamizole to acetic acid-induced writhing in mice. *Folia Pharmacologica Japonica*, 73(8), 907–917. <https://doi.org/10.1254/fpj.73.907>
- [257] Nagaraja, G. K., Prakash, G. K., Kumaraswamy, M. N., Vaidya, V. P., & Mahadevan, K. M. (2006). Synthesis of novel nitrogen containing naphtho[2,1-b]furan derivatives and investigation of their anti microbial activities. *General Papers ARKIVOC*, 15(1), 160–168.
- [258] Naik, H. R. P., Naik, H. S. B., Aravinda, T., & Lamani, D. S. (2010). Synthesis, characterization, and photoactivated DNA cleavage by copper (II)/cobalt (II) mediated macrocyclic complexes. *Nucleosides, Nucleotides and Nucleic Acids*, 29(1), 39–48. <https://doi.org/10.1080/15257770903451496>
- [259] Najafi, Z., Mahdavi, M., Saeedi, M., Karimpour-Razkenari, E., Asatouri, R., Vafadarnejad, F., Moghadam, F. H., Khanavi, M., Sharifzadeh, M., & Akbarzadeh, T. (2017). Novel tacrine-1,2,3-triazole hybrids: In vitro, in vivo biological evaluation and docking study of cholinesterase inhibitors. *European Journal of Medicinal Chemistry*, 125, 1200–1212. <https://doi.org/10.1016/j.ejmech.2016.11.008>

- [260] Nammalwar, B., Muddala, N., Bourne, C., Henry, M., Bourne, P., Bunce, R., Barrow, E., Berlin, K., & Barrow, W. (2014). Synthesis and Biological Evaluation of 2,4-Diaminopyrimidine-Based Antifolate Drugs against *Bacillus anthracis*. *Molecules*, 19(3), 3231–3246. <https://doi.org/10.3390/molecules19033231>
- [261] Nandeshwarappa, B. P., Manjappa, S., & Kishore, B. (2011). A novel approach toward the synthesis of azetidinones derivatives. *Journal of Sulfur Chemistry*, 32(5), 475–481. <https://doi.org/10.1080/17415993.2011.601870>
- [262] Nath, P., Dhumwad, S. D., & Dhumwad, S. D. (2012). Antimicrobial studies of Co(II), Ni(II), Cu(II) and Zn(II) complexes derived from schiff bases of 3-formyl quinoline and 3-hydrazinoquinoxalin-2(1H)one. *Rasayan J Chem*, 5(2), 234–238. <http://www.rasayanjournal.com>
- [263] Nem~ekov/T, E., Rybfir Z', A., Nosfil'ovfi, G., & Alfiildi, J. (1995). Chemical Monthly 7-Substituted 8-Vinyl-1,3-dimethyl-3,7-dihydro-1H-purine-2,6-diones and their Broncholytic Activity. In *Monatshefte fur Chemie* (Vol. 126, Issue 1, pp. 799–804).
- [264] Nikitenko, A. A., Winkley, M. W., Zeldis, J., Kremer, K., Chan, A. W. Y., Strong, H., Jennings, M., Jirkovsky, I., Blum, D., Khafizova, G., Grosu, G. T., & Venkatesan, A. M. (2006). Selective hydrolysis of ethyl 5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole-2-carboxylate and ethyl 5,6-dihydro-4H-pyrrolo[1,2-b]pyrazole-3-carboxylate as a key step in the large-scale synthesis of bicyclic heteroaryl carboxyaldehydes. *Organic Process Research and Development*, 10(4), 712–716. <https://doi.org/10.1021/op050218b>
- [265] Nipate, D. S., Jaspal, S., Shinde, V. N., Rangan, K., & Kumar, A. (2021). TEMPO-Mediated Cross-Dehydrogenative Coupling of Indoles and Imidazo[1,2- a]pyridines with Fluorinated Alcohols. *Organic Letters*, 23(4), 1373–1377. <https://doi.org/10.1021/acs.orglett.1c00031>
- [266] Noureldin, N. A., Richards, J., Kothayer, H., Baraka, M. M., Eladl, S. M., Wootton, M., & Simons, C. (2021). Design, computational studies, synthesis and in vitro antimicrobial evaluation of benzimidazole based thio-oxadiazole and thio-thiadiazole analogues. *BMC Chemistry*, 15(1), 1–19. <https://doi.org/10.1186/s13065-021-00785-8>

- [267] O.A., E. S. M. , E. S. M. A. , Khalil. (1996). Non-steroidal anti-inflammatory agents: synthesis of pyrazolyl pyrazolinyl and pyrimidinyl derivatives of quinolone. *Alexandria Journal of Pharmaceutical Sciences*, 1(10), 43–46.
- [268] Obot, I. B., & Johnson, A. (2012). Ab initio, DFT and TD-DFT electronic absorption spectra investigations on 3,5-diamino-1,2,4-triazole. *Elixir Compound Chemistry*, 43, 6658–6661. <https://www.researchgate.net/publication/261725304>
- [269] Ohtsuka, Y., Uraguchi, D., Yamamoto, K., Tokuhisa, K., & Yamakawa, T. (2012). Syntheses of 2-(trifluoromethyl)-1,3-dicarbonyl compounds through direct trifluoromethylation with CF₃I and their application to fluorinated pyrazoles syntheses. *Tetrahedron*, 68(12), 2636–2649. <https://doi.org/10.1016/j.tet.2012.01.075>
- [270] Okonya, J. F., Hoffman, R. V., & Johnson, M. C. (2002). Synthesis of 2-Oxazolone-4-carboxylates from 3-Nosyloxy- and 3-Bromo-2-ketoesters. *The Journal of Organic Chemistry*, 67(4), 1102–1108. <https://doi.org/10.1021/jo010630z>
- [271] Onatibia-Astibia, A., Franco, R., & Martinez-Pinilla, E. (2017). Health benefits of methylxanthines in neurodegenerative diseases. *Molecular Nutrition & Food Research*, 61(6). <https://doi.org/10.1002/mnfr.201600670>
- [272] Orrego Hernandez, J., Portilla, J., Cobo, J., & Glidewell, C. (2015). Two different products from the reaction of 1-aryl-5-chloro-3-methyl-1H-pyrazole-4-carbaldehyde with cyclohexylamine when the aryl substituent is phenyl or pyridin-2-yl: Hydrogen-bonded sheets versus dimers. *Acta Crystallographica Section C: Structural Chemistry*, 71, 363–368. <https://doi.org/10.1107/S2053229615006403>
- [273] Osman, M., ElSherief, H., & Abdel-Rahman, H. (2018). Synthesis of 1,2,4-triazole derivatives and evaluation of their antioxidant activity. *Journal of Advanced Biomedical and Pharmaceutical Sciences*, 0(0), 0–0. <https://doi.org/10.21608/jabps.2018.3527.1005>
- [274] P Subhashini, N. J., Amanaganti, J., Boddu, L., & Nagarjuna, A. P. (2013). Microwave assisted synthesis and antibacterial studies of (E)-3-(2-Morpholinoquinolin-3-yl)-1-aryl prop-2-en-1-ones. *Available Online Wwww.Jocpr.Com Journal of Chemical and Pharmaceutical Research*, 5(1), 140–147. www.jocpr.com

- [275] P. Venkatesh. (2011). Synthesis, Characterisation And Anti Microbial Activity Of Various Schiff Base Complexes Of Zinc (II) And Copper (II) Ions. *Asian Journal of Pharmaceutical and Health Sciences*, 1(1), 8–11.
- [276] Pachuta-Stec, A., Nowak, R., Pietrzak, W., & Pitucha, M. (2019). Synthesis and Antioxidant Activity of New Norcantharidin Analogs. *Chemistry and Biodiversity*, 16(4). <https://doi.org/10.1002/cbdv.201800673>
- [277] Padmini, thatavarthi, & kamal, bigala raj. (2019). Synthesis, Anti-inflammatory, Analgesic and Antipyretic Activity of Novel 1,3,5-Trisubstituted Pyrazole Derivatives. *Asian Journal of Chemistry*, 31(6), 1225–1229. <https://doi.org/10.14233/ajchem.2019.21781>
- [278] Palaska, E., lay, G. S., Kelicen, P., Tug, N., Durlu, ba, & lçin Altinok, G. (2002). Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides. *Il Farmaco*, 57, 101–107. www.elsevier.com/locate/farmac
- [279] Pandey, A., Rajavel, R., Chandraker, S., & Dash, D. (2012). Synthesis of Schiff Bases of 2-amino-5-aryl-1,3,4-thiadiazole and Its Analgesic, Anti-Inflammatory and Anti-Bacterial Activity. *E-Journal of Chemistry*, 9(4), 2524–2531. <http://www.ejchem.net>
- [280] Pandey, V. K., Tusi, S., Misra, R., & Shukla, & R. (2010). A chemical strategy for the construction of quinoline isoquinoline core units. In *Indian Journal of Chemistry* (Vol. 49).
- [281] Pandit, A. B., Savant, M. M., & Ladva, K. D. (2018). An Efficient One-Pot Synthesis of Highly Substituted Pyridone Derivatives and Their Antimicrobial and Antifungal Activity. *Journal of Heterocyclic Chemistry*, 55(4), 983–987. <https://doi.org/10.1002/jhet.3128>
- [282] Pandit, U., & Dodiya, A. (2013). Synthesis and antitubercular activity of novel pyrazole–quinazolinone hybrid analogs. *Medicinal Chemistry Research*, 22(7), 3364–3371. <https://doi.org/10.1007/s00044-012-0351-0>
- [283] Parmentier-Batteur, S., OBrien, J. A., Doran, S., Nguyen, S. J., Flick, R. B., Uslander, J. M., Chen, H., Finger, E. N., Williams, T. M., Jacobson, M. A., & Hutson, P. H. (2012). Differential effects of the mGluR5 positive allosteric modulator CDPPB in the cortex and striatum following repeated administration. *Neuropharmacology*, 62(3), 1453–1460. <https://doi.org/10.1016/j.neuropharm.2010.11.013>

- [284] Parrick, J., Mehta, L. K., & Hodgkiss, R. J. (1993). The synthesis of a potential anti-cancer agent containing the caffeine and 1,2,4-benzotriazine moieties. *Journal of Heterocyclic Chemistry*, 30(2), 323–327. <https://doi.org/10.1002/jhet.5570300205>
- [285] Paruch, K., Popiolek, lukasz, & Wujec, M. (2020). Antimicrobial and antiprotozoal activity of 3-acetyl-2,5-disubstituted-1,3,4-oxadiazolines: a review. *Medicinal Chemistry Research*, 29(1), 1–16. <https://doi.org/10.1007/s00044-019-02463-w>
- [286] Patel, K. D. , M. B. D. , & D. K. R. (2002). Synthesis and antimicrobial activity of 1, 2, 4-triazoles . *Journal of the Indian Chemical Society*, 12(79), 964-965.
- [287] Pathak, R. B., Chovatia, P. T., & Parekh, H. H. (2012). Synthesis, antitubercular and antimicrobial evaluation of 3-(4-chlorophenyl)-4-substituted pyrazole derivatives. *Bioorganic & Medicinal Chemistry Letters*, 22(15), 5129–5133. <https://doi.org/10.1016/j.bmcl.2012.05.063>
- [288] Patin, A., & Belmont, P. (2005). A New Route to Acridines: Pauson-Khand Reaction on Quinoline-Bearing 1-En-7-yne Leading to Novel Tetrahydrocyclopenta[c]acridine-2,5-diones. *Synthesis*, 14(14), 2400–2406. <https://doi.org/10.1055/s-2005-870016>
- [289] Phatak, P. S., Sathe, B. P., Dhumal, S. T., Rehman, N. N. M. A., Dixit, P. P., Khedkar, V. M., & Haval, K. P. (2019). Synthesis, Antimicrobial Evaluation, and Docking Studies of Substituted Acetylphenoxymethyl-triazolyl-N-phenylacetamides. *Journal of Heterocyclic Chemistry*, 56(7), 1928–1938. <https://doi.org/10.1002/jhet.3568>
- [290] Pillai, R. R., Karrouchi, K., Fettach, S., Armarković, S., Armarković, S. J., Brik, Y., Taoufik, J., Radi, S., El Abbes Faouzi, M., & Ansar, M. (2019). Synthesis, spectroscopic characterization, reactive properties by DFT calculations, molecular dynamics simulations and biological evaluation of Schiff bases tethered 1,2,4-triazole and pyrazole rings. *Journal of Molecular Structure*, 1177, 47–54. <https://doi.org/10.1016/j.molstruc.2018.09.037>
- [291] Pokalwar, R. U., Hangarge, R. V., Madje, B. R., Ware, M. N., & Shingare, M. S. (2008). Simple and high yielding synthesis of new α -aminophosphonates from imines. *Phosphorus, Sulfur and Silicon and the Related Elements*, 183(6), 1461–1470. <https://doi.org/10.1080/10426500701681532>

- [292] Pokalwar, R. U., Hangarge, R. V., Maske, P. V., & Shingare, M. S. (2006). Synthesis and antibacterial activities of α -hydroxyphosphonates and α -acetyloxyphosphonates derived from 2-chloroquinoline-3-carbaldehyde. *Arkivoc*, 2006(11), 196–204. <https://doi.org/10.3998/ark.5550190.0007.b20>
- [293] Ponnala, S., Kiran Kumar, S. T. V. S., Bhat, B. A., & Prasad Sahu, D. (2005). Synthesis of Bridgehead Nitrogen Heterocycles on a Solid Surface. *Synthetic Communications*, 35(7), 901–906. <https://doi.org/10.1081/SCC-200051674>
- [294] Ponnala, S., & Prasad Sahu, D. (2006). Iodine-mediated synthesis of 2-arylbenzoxazoles, 2-arylbenzimidazoles, and 1,3,5-trisubstituted pyrazoles. *Synthetic Communications*, 36(15), 2189–2194. <https://doi.org/10.1080/00397910600638879>
- [295] Potts, K. T. (1960). *The chemistry of 1,2,4-triazole*.
- [296] Prajapati, D., & Borah, K. J. (2007). The tert-amino effect in heterocyclic chemistry: Synthesis of new fused pyrazolinoquinolizine and 1,4-oxazinopyrazoline derivatives. *Beilstein Journal of Organic Chemistry*, 3. <https://doi.org/10.1186/1860-5397-3-43>
- [297] Pramod Kumar Dubey and A.Naidu and K. Shiva Kumar. (2008). Microwave assisted, solvent-free synthesis of quinolinyl-1,2,4-triazolo[4,3-a]quinoxalines using IBD. *Indian Journal of Organic Chemistry*, 4.
- [298] Praveen, S., Kumar, A., Rabiya Parveen, B., & Pal, H. (2017). International Journal of Pharmaceutical Chemistry Synthesis and biological evaluation of some new Schiff base 1,2,4-triazole derivatives. *International Journal of Pharmaceutical Chemistry*, 7(6), 95–99. <https://doi.org/10.7439/ijpc>
- [299] Punia, S., Verma, V., Kumar, D., Kumar, A., & Deswal, L. (2021). Facile synthesis, antimicrobial evaluation and molecular docking studies of pyrazole-imidazole-triazole hybrids. *Journal of Molecular Structure*, 1223. <https://doi.org/10.1016/j.molstruc.2020.129216>
- [300] Puthiaraj, P., Ramu, A., & Pitchumani, K. (2014). Copper-based metal-organic frameworks as reusable heterogeneous catalysts for the one-pot syntheses of imidazo[1,2-a]pyridines. *Asian Journal of Organic Chemistry*, 3(7), 784–791. <https://doi.org/10.1002/ajoc.201402019>
- [301] Quattrini, L., Gelardi, E. L. M., Coviello, V., Sartini, S., Ferraris, D. M., Mori, M., Nakano, I., Garavaglia, S., & La Motta, C. (2020). Imidazo[1,2- a]pyridine Derivatives as Aldehyde Dehydrogenase Inhibitors: Novel Chemotypes to Target

- Glioblastoma Stem Cells. *Journal of Medicinal Chemistry*, 63(9), 4603–4616. <https://doi.org/10.1021/acs.jmedchem.9b01910>
- [302] R, K., & Bodke, Y. D. (2020). Synthesis, analgesic and anti-inflammatory activity of benzofuran pyrazole heterocycles. *Chemical Data Collections*, 28. <https://doi.org/10.1016/j.cdc.2020.100453>
- [303] Rabe, K., Magnussen, H., & Dent, G. (1995). Theophylline and selective PDE inhibitors as bronchodilators and smooth muscle relaxants. *European Respiratory Journal*, 8(4), 637–642. <https://doi.org/10.1183/09031936.95.08040637>
- [304] Raghu Prasad, M., & Pran Kishore, D. (2007). Multistep, Microwave Assisted, Solvent Free Synthesis and Antibacterial Activity of 6-Substituted-2,3,4-trihydropyrimido[1,2-c]9,10,11,12-tetrahydrobenzo[b]thieno[3,2-e]pyrimidines. *Chemical and Pharmaceutical Bulletin*, 55(5), 776–779. <https://doi.org/10.1248/cpb.55.776>
- [305] Raghu Prasad, M., Rao, A. R. R., Shanthan Rao, P., & Subramanian Rajan, K. (2002). Microwave-Assisted Synthesis of Novel 5-Substituted-2,3-dihydroimidazo[1,2-c]thieno[3,2-e]pyrimidines. *ChemInform*, 33(8). <https://doi.org/10.1002/chin.200208165>
- [306] Rajak, H., Agarawal, A., Parmar, P., Thakur, B. S., Veerasamy, R., Sharma, P. C., & Kharya, M. D. (2011). 2,5-Disubstituted-1,3,4-oxadiazoles/thiadiazole as surface recognition moiety: Design and synthesis of novel hydroxamic acid based histone deacetylase inhibitors. *Bioorganic and Medicinal Chemistry Letters*, 21(19), 5735–5738. <https://doi.org/10.1016/j.bmcl.2011.08.022>
- [307] Ramadan, E. S., Sharshira, E. M., El Sökkary, R. I., & Morsy, N. (2018). Synthesis and antimicrobial evaluation of some heterocyclic compounds from 3-Aryl-1-phenyl-1H-pyrazole-4-carbaldehydes. *Zeitschrift Fur Naturforschung - Section B Journal of Chemical Sciences*, 73(6), 389–397. <https://doi.org/10.1515/znb-2018-0009>
- [308] Rao, G. , R. S. , & A. M. (2000). Synthesis and antimicrobial acitivity of some 5-phenyl-4-substituted amino-3-mercapto-(4H)-1,2,4-triazoles. *Indian Journal of Pharmaceutical Sciences*, 6(62), 475-477.
- [309] Rashid, M., Husain, A., & Mishra, R. (2012). Synthesis of benzimidazoles bearing oxadiazole nucleus as anticancer agents. *European Journal of Medicinal Chemistry*, 54, 855–866. <https://doi.org/10.1016/j.ejmech.2012.04.027>

- [310] Rathore, A., Sudhakar, R., Ahsan, M. J., Ali, A., Subbarao, N., Jadav, S. S., Umar, S., & Yar, M. S. (2017). In vivo anti-inflammatory activity and docking study of newly synthesized benzimidazole derivatives bearing oxadiazole and morpholine rings. *Bioorganic Chemistry*, 70, 107–117. <https://doi.org/10.1016/j.bioorg.2016.11.014>
- [311] Raundal, H. N., Jadhav, R. P., Patil, A. A., & Bobade, V. D. (2014a). Synthesis and antimicrobial studies of 2-(5-substituted)-1, 3, 4-oxadiazole-2-yl)-H-imidazo [1, 2, α] pyridine derivatives. Available Online *Www.Jocpr.Com Journal of Chemical and Pharmaceutical Research*, 6(7), 102–108. www.jocpr.com
- [312] Raundal, H. N., Jadhav, R. P., Patil, A. A., & Bobade, V. D. (2014b). Synthesis and antimicrobial studies of 2-(5-substituted)-1, 3, 4-oxadiazole-2-yl)-H-imidazo[1, 2, α] pyridine derivatives. *Journal of Chemical and Pharmaceutical Research*, 6(7), 102–108. www.jocpr.com
- [313] Raydan, D., Friaes, S., Viduedo, N., Santos, A. S., Gomes, C. S. B., Royo, B., & Marques, M. M. B. (2022). Manganese-Catalyzed Synthesis of Imines from Primary Alcohols and (Hetero)Aromatic Amines. *Synlett*, 33(13), 1290–1294. <https://doi.org/10.1055/a-1828-1678>
- [314] Reddyrajula, R., & Dalimba, U. (2020). The bioisosteric modification of pyrazinamide derivatives led to potent antitubercular agents: Synthesis via click approach and molecular docking of pyrazine-1,2,3-triazoles. *Bioorganic and Medicinal Chemistry Letters*, 30(2). <https://doi.org/10.1016/j.bmcl.2019.126846>
- [315] Reddyrajula, R., & Dalimba, U. K. (2019). Structural modification of zolpidem led to potent antimicrobial activity in imidazo[1,2- A] pyridine/pyrimidine-1,2,3-triazoles. *New Journal of Chemistry*, 43(41), 16281–16299. <https://doi.org/10.1039/c9nj03462e>
- [316] Romero, A. H. (2016). Modified procedure for the synthesis of 2-chloroquinoline-3-carbaldehydes using phosphorus pentachloride. *Synthetic Communications*, 46(3), 287–291. <https://doi.org/10.1080/00397911.2015.1135956>
- [317] Rostovtsev, V. V., Green, L. G., Fokin, V. V., Barry Sharpless, K., A C Coolen, H. K., N M van Leeuwen, P. W., & M Nolte, R. J. (2002). Harrowfield in Calixarenes. In *J. Chem. Soc. Chem. Commun* (Vol. 114). Kluwer Academic Publishers. <http://www.angewandte.org>

- [318] Rozin, P., Levine, E., & Stoess, C. (1991). Chocolate craving and liking. *Appetite*, 17(3), 199–212. [https://doi.org/10.1016/0195-6663\(91\)90022-K](https://doi.org/10.1016/0195-6663(91)90022-K)
- [319] Ruddaraju, R. R., Murugulla, A. C., Kotla, R., Tirumalasetty, M. C. B., Wudayagiri, R., Donthabakthuni, S., & Maraju, R. (2017). Design, synthesis, anticancer activity and docking studies of theophylline containing 1,2,3-triazoles with variant amide derivatives. *MedChemComm*, 8(1), 176–183. <https://doi.org/10.1039/C6MD00479B>
- [320] Russo, C. M., Adhikari, A. A., Wallach, D. R., Fernandes, S., Balch, A. N., Kerr, W. G., & Chisholm, J. D. (2015). Synthesis and initial evaluation of quinoline-based inhibitors of the SH2-containing inositol 5'-phosphatase (SHIP). *Bioorganic & Medicinal Chemistry Letters*, 25(22), 5344–5348. <https://doi.org/10.1016/j.bmcl.2015.09.034>
- [321] Saaïd, F. H., Dawood, S. J., & Marbeen, B. H. (2018). Synthesis and Characterization of 1,3,4- oxadiazole Derivatives using an Ultrasonic Technique. *Al-Mustansiriyah Journal of Science*, 29(2), 101–105. <https://doi.org/10.23851/mjs.v29i2.183>
- [322] Sahoo, S., Patwari, P. K., Kumar, M. C., & Setty, Cm. (2013). Synthesis and Biological Activity of Certain Mannich Bases Derivatives from 1, 2, 4-Triazoles. *Iranian Journal of Pharmaceutical Sciences*, 2013(9), 51–60. www.ijps.ir
- [323] Sahu, J. K., Ganguly, S., & Kaushik, A. (2014). Synthesis of some novel heterocyclic 1,2,4-triazolo [3,4-b][1,3,4] thiadiazole derivatives as possible antimicrobial agents. *Journal of Applied Pharmaceutical Science*, 4(2), 81–86. <https://doi.org/10.7324/JAPS.2014.40214>
- [324] Saini, L., kumar Yadav, A., Yadav, P., Kumar Gupta, S., & Srivastava, S. (2015). Synthesis, Characterization and biological activity of some novel pyrazole derivatives. *World Journal of Pharmaceutical Research*, 12(3), 744–755. <https://doi.org/10.20959/wjpr20233-27045>
- [325] Saini, R. P., Kumar, V., Gupta, A. K., & Gupta, G. K. (2014). Synthesis, characterization, and antibacterial activity of a novel heterocyclic Schiff's base and its metal complexes of first transition series. *Medicinal Chemistry Research*, 23(2), 690–698. <https://doi.org/10.1007/s00044-013-0657-6>

- [326] Saleh, N. M., El-Gazzar, M. G., Aly, H. M., & Othman, R. A. (2020). Novel Anticancer Fused Pyrazole Derivatives as EGFR and VEGFR-2 Dual TK Inhibitors. *Frontiers in Chemistry*, 7. <https://doi.org/10.3389/fchem.2019.00917>
- [327] Salih, N. A., Abd, H., & Ibraheem, E.-L. (2008). A New Derivatives of Benzodiazepine, Imidazole, Isatin, Maleimide, Pyrimidine and 1,2,4-Triazole: Synthesis and Characterization. *Um-Salama Science Journal*, 5(2).
- [328] Sathe, B. S. , J. E. , J. V. A. , & S. (2011). Synthesis characterization and anti-inflammatory evaluation of new fluorobenzothiazole schiff's bases. *Int J Pharm Res Dev*, 3(3), 164–169.
- [329] Schiff, H. (1864). Mittheilungen aus dem Universitätslaboratorium in Pisa: Eine neue Reihe organischer Basen. *Justus Liebigs Annalen Der Chemie*, 131(1), 118–119. <https://doi.org/10.1002/jlac.18641310113>
- [330] Sekar, M., & Rajendra Prasad, K. J. (1998). Synthesis of Some Novel 2-Oxo-pyrano(2,3-b)-and 2-Oxo-pyrido(2,3-b)quinoline Derivatives as Potential Antimalarial, Diuretic, Clastogenic and Antimicrobial Agents. *J. Chem. T Echnol. Biotechnol*, 72, 50–54.
- [331] Selvi, S. T., Nadaraj, V., Mohan, S., Sasi, R., & Hema, M. (2006). Solvent free microwave synthesis and evaluation of antimicrobial activity of pyrimido[4,5-b]- and pyrazolo[3,4-b]quinolines. *Bioorganic & Medicinal Chemistry*, 14(11), 3896–3903. <https://doi.org/10.1016/j.bmc.2006.01.048>
- [332] Serafini, M., Cordero-Sanchez, C., Di Paola, R., Bhela, I. P., Aprile, S., Purghè, B., Fusco, R., Cuzzocrea, S., Genazzani, A. A., Riva, B., & Pirali, T. (2020). Store-Operated Calcium Entry as a Therapeutic Target in Acute Pancreatitis: Discovery and Development of Drug-Like SOCE Inhibitors. *Journal of Medicinal Chemistry*, 63(23), 14761–14779. <https://doi.org/10.1021/acs.jmedchem.0c01305>
- [333] Shapranova, N. I., Somin, I. N., & Geterotsiklicheskikh, K. (1970). The synthesis of 1-methylpyrazolealdehydes and 1-methyl-2-pyrazolinealdehydes and their acetals. *Khimiya Geterotsiklicheskikh Soedinenii*, 6(3), 404–406.
- [334] Shawali, A. S., Mosselhi, M. A. N., & Farghaly, T. A. (2007). Synthesis and Tautomeric Structure of 2-arylaazo-4H -imidazo[2,1-b][1,3,4]thiadiazines. *Journal of Chemical Research*, 2007(8), 479–483. <https://doi.org/10.3184/030823407X237885>

- [335] Sheeja, K., Beula Priyanka, G., Arul Salomon, T., Bhagya Lakshmi, S., Vijaya Kumar, P., & Suchitra, M. (2013). Design, synthesis, characterization and biological activity studies on coumarin carbohydrazide containing quinoline derivatives. *Asian Journal of Pharmaceutical Analysis and Medicinal Chemistry*, 1(1), 39–47. www.uptodateresearchpublication.com
- [336] Sheldrick, G. M. (2015). Crystal structure refinement with SHELXL. *Acta Crystallographica Section C: Structural Chemistry*, 71, 3–8. <https://doi.org/10.1107/S2053229614024218>
- [337] Shelke RN. (2017). *Synthesis of 2-((substituted)-2-chloroquinolin-3-yl)-3-((substituted) phenyl) thiazolidin-4-one with β -cyclodextrin-SO₃H Catalyst Under Solvent-free Condition.*
- [338] Sheremet, E. A., Tomanov, R. I., Trukhin, E. V., & Berestovitskaya, V. M. (2004). Synthesis of 4-Aryl-5-nitro-1,2,3-triazoles. *Russian Journal of Organic Chemistry*, 40(4), 594–595. <https://doi.org/10.1023/B:RUJO.0000036090.61432.18>
- [339] Shin, J. A., Jung, H., & Lim, Y. G. (2020). Competitive CuAAC reaction between hydrophobic and hydrophilic alkynes with azides in water. *ChemistrySelect*, 5(40), 12371–12376. <https://doi.org/10.1002/slct.202002792>
- [340] Shneine, J., Alaraji, Y., Shneine, J. K., & Alaraji, Y. H. (2016). Chemistry of 1, 2, 4-Triazole: A Review Article. *International Journal of Science and Research*, 5(3). <https://www.researchgate.net/publication/312021996>
- [341] Shyam Maurya, S., Priya Gosain, T., Kidwai, S., Singh, R., Diwan, &, & Rawat, S. (2019). Synthesis of 1,3,4-oxadiazole and imidazo[1,2-a]pyridine based molecular hybrids and their in vitro antituberculosis and cytotoxicity studies. *Indian Journal of Chemistry*, 58, 1005–1018.
- [342] Singh, P., Swain, B., Thacker, P. S., Sigalapalli, D. K., Purnachander Yadav, P., Angeli, A., Supuran, C. T., & Arifuddin, M. (2020). Synthesis and carbonic anhydrase inhibition studies of sulfonamide based indole-1,2,3-triazole chalcone hybrids. *Bioorganic Chemistry*, 99. <https://doi.org/10.1016/j.bioorg.2020.103839>
- [343] Singh, R., Kashaw, S. K., Mishra, V. K., Mishra, M., Rajoriya, V., & Kashaw, V. (2018). Design and Synthesis of New Bioactive 1,2,4-Triazoles, Potential Antitubercular and Antimicrobial Agents. *Indian Journal of Pharmaceutical Sciences*, 80(01). <https://doi.org/10.4172/pharmaceutical-sciences.1000328>

- [344] Smeyers, Y. (1989). study comparative CNDO/2 des determinants moleculaires de liactivite agoniste de recepteur-H2 dans histamine et le betazol. *European Journal of Medicinal Chemistry*, 24(4), 411–414. [https://doi.org/10.1016/0223-5234\(89\)90085-8](https://doi.org/10.1016/0223-5234(89)90085-8)
- [345] Somani, R. R., & Shirodkar, P. Y. (2009). Oxadiazole: A biologically important heterocycle. In *Pharma Chemica* (Vol. 1, Issue 1).
- [346] Sondhi, S. M., Kumar, S., Kumar, N., & Roy, P. (2012). Synthesis anti-inflammatory and anticancer activity evaluation of some pyrazole and oxadiazole derivatives. *Medicinal Chemistry Research*, 21(10), 3043–3052. <https://doi.org/10.1007/s00044-011-9850-7>
- [347] Sondhi, S. M., Singh, N., Kumar, A., Lozach, O., & Meijer, L. (2006). Synthesis, anti-inflammatory, analgesic and kinase (CDK-1, CDK-5 and GSK-3) inhibition activity evaluation of benzimidazole/benzoxazole derivatives and some Schiff's bases. *Bioorganic & Medicinal Chemistry*, 14(11), 3758–3765. <https://doi.org/10.1016/j.bmc.2006.01.054>
- [348] Spano, V., Parrino, B., Carbone, A., Montalbano, A., Salvador, A., Brun, P., Vedaldi, D., Diana, P., Cirrincione, G., & Barraja, P. (2015). Pyrazolo[3,4-h]quinolines promising photosensitizing agents in the treatment of cancer. *European Journal of Medicinal Chemistry*, 102, 334–351. <https://doi.org/10.1016/j.ejmech.2015.08.003>
- [349] Spitz, I. M., Novis, B. H., Ebert, R., Trestian, S., LeRoith, D., & Creutzfeldt, W. (1982). Betazole-induced GIP secretion is not mediated by gastric HCl. *Metabolism*, 31(4), 380–382. [https://doi.org/10.1016/0026-0495\(82\)90114-7](https://doi.org/10.1016/0026-0495(82)90114-7)
- [350] Srivastava, A., Singh, M. K., & Singh, R. M. (2006). Pyrazolo-fused quinoline analogues: Synthesis of 1H-pyrazolo [3, 4-b] quinolines and 3-amino-1H-pyrazolo [3, 4-b] quinolines from 3-formyl and 3-cyano-2-chloroquinolines. *Indian Journal of Chemistry*, 458, 292–296.
- [351] Stasyuk, A. J., Banasiewicz, M., Cyrański, M. K., & Gryko, D. T. (2012). Imidazo[1,2-a]pyridines susceptible to excited state intramolecular proton transfer: One-pot synthesis via an Ortoleva-King reaction. *Journal of Organic Chemistry*, 77(13), 5552–5558. <https://doi.org/10.1021/jo300643w>
- [352] Steinbach, G., Lynch, P. M., Phillips, R. K. S., Wallace, M. H., Hawk, E., Gordon, G. B., Wakabayashi, N., Saunders, B., Shen, Y., Fujimura, T., Su, L.-K., Levin, B., Godio, L., Patterson, S., Rodriguez-Bigas, M. A., Jester, S. L., King, K. L.,

- Schumacher, M., Abbruzzese, J., ... Kelloff, G. (2000). The Effect of Celecoxib, a Cyclooxygenase-2 Inhibitor, in Familial Adenomatous Polyposis. *New England Journal of Medicine*, 342(26), 1946–1952. <https://doi.org/10.1056/NEJM200006293422603>
- [353] Sumangala, V., Poojary, B., Chidananda, N., Fernandes, J., & Kumari, N. S. (2010). Synthesis and antimicrobial activity of 1,2,3-triazoles containing quinoline moiety. *Archives of Pharmacal Research*, 33(12), 1911–1918. <https://doi.org/10.1007/s12272-010-1204-3>
- [354] Sundaramoorthy, R., Vadivelu, M., Thirumoorthy, K., Karthikeyan, K., & Praveen, C. (2023). Step-Economical Mechanosynthesis of Hybrid Azoles: Deciphering Their π -Orbital and Pharmacological Characteristics. *ChemMedChem*, 18(13). <https://doi.org/10.1002/cmdc.202300008>
- [355] Szabo, G., Fischer, J., Kis-Varga, A., & Gyires, K. (2008). New Celecoxib Derivatives as Anti-Inflammatory Agents. *Journal of Medicinal Chemistry*, 51(1), 142–147. <https://doi.org/10.1021/jm070821f>
- [356] Tan, W., Li, Q., Li, W., Dong, F., & Guo, Z. (2016). Synthesis and antioxidant property of novel 1,2,3-triazole-linked starch derivatives via “click chemistry.” *International Journal of Biological Macromolecules*, 82, 404–410. <https://doi.org/10.1016/j.ijbiomac.2015.10.007>
- [357] Tekale, A. S., Mukhedker, S. S., & Shaikh, S. A. L. (2015). A highly efficient synthesis of 2-chloro-3-formyl-8-methyl quinoline: Vilsmeier-haack reagent. ~ 42 ~ *International Journal of Chemical Studies*, 2(6), 42–45.
- [358] Terrett, N. K. (2013). Combinatorial Chemistry Online. *Combinatorial Chemistry - an Online Journal*, 15(4), 13–15. <https://doi.org/10.1016/j.comche.2013.03.001>
- [359] Thakare, P. P., Walunj, Y., Chavan, A., Bobade, V. D., Sarkar, D., & Mhaske, P. C. (2020). Synthesis and antimycobacterial screening of new 4-(4-(1-benzyl-1H-1,2,3-triazol-4-yl)-1-phenyl-1H-pyrazol-3-yl)quinoline derivatives. *Journal of Heterocyclic Chemistry*, 57(11), 3918–3929. <https://doi.org/10.1002/jhet.4101>
- [360] Thakor, T., & Savjani, J. (2014). Synthesis and Cell line study of Pyrazole Substituted Coumarin Derivatives. In *International Journal of PharmTech Research CODEN (USA): IJPRIF ISSN* (Vol. 6, Issue 4, pp. 1397–1406).

- [361] Tornøe, C. W., Christensen, C., & Meldal, M. (2002). Peptidotriazoles on Solid Phase: [1,2,3]-Triazoles by Regiospecific Copper(I)-Catalyzed 1,3-Dipolar Cycloadditions of Terminal Alkynes to Azides. *The Journal of Organic Chemistry*, 67(9), 3057–3064. <https://doi.org/10.1021/jo0011148j>
- [362] Tozkoparan, B., Kupeli, E., Yesilada, E., & Ertan, M. (2007). Preparation of 5-aryl-3-alkylthio-1,2,4-triazoles and corresponding sulfones with antiinflammatory-analgesic activity. *Bioorganic and Medicinal Chemistry*, 15(4), 1808–1814. <https://doi.org/10.1016/j.bmc.2006.11.029>
- [363] Trivedi, H. D., Joshi, V. B., & Patel, B. Y. (2022). Pyrazole Bearing Pyrimidine Analogues as the Privileged Scaffolds in Antimicrobial Drug Discovery: A Review. *Analytical Chemistry Letters*, 12(2), 147–173. <https://doi.org/10.1080/22297928.2021.1910565>
- [364] Tsutomu, K., & Toshitaka, N. (1978). Effects of 1,3-diphenyl-5-(2-dimethylaminopropionamide)-pyrazole[difenamizole] on a conditioned avoidance response. *Neuropharmacology*, 17(4–5), 249–256. [https://doi.org/10.1016/0028-3908\(78\)90108-9](https://doi.org/10.1016/0028-3908(78)90108-9)
- [365] Tzanetou, E., Liekens, S., Kasiotis, K. M., Fokialakis, N., & Haroutounian, S. A. (2012). Novel Pyrazole and Indazole Derivatives: Synthesis and Evaluation of Their Anti-Proliferative and Anti-Angiogenic Activities. *Archive of Pharmacy*, 345(10), 804–811. <https://doi.org/10.1002/ardp.201200057>
- [366] Ulloora, S., Shabaraya, R., & Adhikari, A. V. (2013). Facile synthesis of new imidazo[1,2-a]pyridines carrying 1,2,3-triazoles via click chemistry and their antiepileptic studies. *Bioorganic & Medicinal Chemistry Letters*, 23(11), 3368–3372. <https://doi.org/10.1016/j.bmcl.2013.03.086>
- [367] Uslander, J. M., Parmentier-Batteur, S., Flick, R. B., Surles, N. O., Lam, J. S. H., McNaughton, C. H., Jacobson, M. A., & Hutson, P. H. (2009). Dose-dependent effect of CDPPB, the mGluR5 positive allosteric modulator, on recognition memory is associated with GluR1 and CREB phosphorylation in the prefrontal cortex and hippocampus. *Neuropharmacology*, 57(5–6), 531–538. <https://doi.org/10.1016/j.neuropharm.2009.07.022>

- [368] Vandekerckhove, S., & D'hooghe, M. (2015). Quinoline-based antimalarial hybrid compounds. *Bioorganic & Medicinal Chemistry*, 23(16), 5098–5119. <https://doi.org/10.1016/j.bmc.2014.12.018>
- [369] Vandekerckhove, S., Van Herreweghe, S., Willems, J., Danneels, B., Desmet, T., de Kock, C., Smith, P. J., Chibale, K., & D'hooghe, M. (2015). Synthesis of functionalized 3-, 5-, 6- and 8-aminoquinolines via intermediate (3-pyrrolin-1-yl)- and (2-oxopyrrolidin-1-yl)quinolines and evaluation of their antiplasmodial and antifungal activity. *European Journal of Medicinal Chemistry*, 92(1), 91–102. <https://doi.org/10.1016/j.ejmech.2014.12.020>
- [370] Vardan, S., Smulyan, H., Mookherjee, S., Eich, R., Syracuse, N. Y., & Vardan, S. (1983). Effects of tiodazosin, a new antihypertensive, hemodynamics and clinical variables. *Clinical Pharmacology and Therapeutics*, 34(3), 290–297. <https://doi.org/https://doi.org/10.1038/clpt.1983.170>
- [371] Visagaperumal, D., Jaya Kumar, R., Vijayaraj, R., & Anbalagan, N. (2009). Microwave induced synthesis of some new substituted-1,3-thiazolidin-4-ones for their potent antimicrobial and antitubercular activities. *International Journal of Chem Tech Research*, 1(4), 1048–1051.
- [372] Vitaku, E., Smith, D. T., & Njardarson, J. T. (2014). Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *Journal of Medicinal Chemistry*, 57(24), 10257–10274. <https://doi.org/10.1021/jm501100b>
- [373] Wall, M. E., Wani, M. C., Cook, C. E., Palmer, K. H., McPhail, A. T., & Sim, G. A. (1966). Plant Antitumor Agents. I. The Isolation and Structure of Camptothecin, a Novel Alkaloidal Leukemia and Tumor Inhibitor from *Camptotheca acuminata*. *Journal of the American Chemical Society*, 88(16), 3888–3890. <https://doi.org/10.1021/ja00968a057>
- [374] Wang, G.-B., Wang, L.-F., Li, C.-Z., Sun, J., Zhou, G.-M., & Yang, D.-C. (2012). A facile and efficient method for the selective deacylation of N-arylacetamides and 2-chloro-N-arylacetamides catalyzed by SOCl₂. *Research on Chemical Intermediates*, 38(1), 77–89. <https://doi.org/10.1007/s11164-011-0327-6>

- [375] Wang, M., Zhang, J., Liu, J., Xu, C., & Ju, H. (2002). Intramolecular energy and charge transfer in 5-(9-anthryl)-3-(4-nitrophenyl)-1-phenyl-2-pyrazoline. *Journal of Luminescence*, 99(1), 79–83. [https://doi.org/10.1016/S0022-2313\(01\)00204-6](https://doi.org/10.1016/S0022-2313(01)00204-6)
- [376] Wang, X. L., Wan, K., & Zhou, C. H. (2010). Synthesis of novel sulfanilamide-derived 1,2,3-triazoles and their evaluation for antibacterial and antifungal activities. *European Journal of Medicinal Chemistry*, 45(10), 4631–4639. <https://doi.org/10.1016/j.ejmech.2010.07.031>
- [377] Watzke, A., Köhn, M., Gutierrez-Rodriguez, M., Wacker, R., Schröder, H., Breinbauer, R., Kuhlmann, J., Alexandrov, K., Niemeyer, C. M., Goody, R. S., & Waldmann, H. (2006). Site-Selective Protein Immobilization by Staudinger Ligation. *Angewandte Chemie International Edition*, 45(9), 1408–1412. <https://doi.org/10.1002/anie.200502057>
- [378] Wazalwar, S. S., Banpurkar, A. R., & Perdih, F. (2017). Aqueous phase synthesis, crystal structure and biological study of isoxazole extensions of pyrazole-4-carbaldehyde derivatives. *Journal of Molecular Structure*, 1150(1), 258–267. <https://doi.org/10.1016/j.molstruc.2017.08.094>
- [379] Wazalwar, S. S., Banpurkar, A. R., & Perdih, F. (2018). Synthesis, Characterization, Molecular Docking Studies and Anticancer Activity of Schiff Bases Derived from 3-(Substituted phenyl)-1-phenyl-1H-pyrazole-4-carbaldehyde and 2-Aminophenol. *Journal of Chemical Crystallography*, 48(4), 185–199. <https://doi.org/10.1007/s10870-018-0727-1>
- [380] Wei, D., Li, N., Lu, G., & Yao, K. (2006). Synthesis, catalytic and biological activity of novel dinuclear copper complex with Schiff base. *Science in China Series B*, 49(3), 225–229. <https://doi.org/10.1007/s11426-006-0225-8>
- [381] Wojcik-Pszczola, K., Jankowska, A., Slusarczyk, M., Jakiela, B., Plutecka, H., Pocięcha, K., Swierczek, A., Popiol, J., Koczurkiewicz-Adamczyk, P., Wyska, E., Pekala, E., Gosens, R., & Chlon-Rzepa, G. (2021). Synthesis and in vitro evaluation of anti-inflammatory, antioxidant, and anti-fibrotic effects of new 8-aminopurine-2,6-dione-based phosphodiesterase inhibitors as promising anti-asthmatic agents. *Bioorganic Chemistry*, 117(1), 1–17. <https://doi.org/10.1016/j.bioorg.2021.105409>

- [382] Yamakawa, T. (2002). In vitro and in vivo antibacterial activity of T-3912, a novel non-fluorinated topical quinolone. *Journal of Antimicrobial Chemotherapy*, 49(3), 455–465. <https://doi.org/10.1093/jac/49.3.455>
- [383] Yassin, F. A. (2010). Novel pyrazolyl pyridazine derivatives likely to possess anti-inflammatory activity. *Journal of Microbiology and Antimicrobials*, 2(7), 93–99. <http://www.academicjournals.org/JMA>
- [384] Ye, J., Mao, L., Xie, L., Zhang, R., Liu, Y., Peng, L., Yang, J., Li, Q., & Yuan, M. (2021). Discovery of a Series of Theophylline Derivatives Containing 1,2,3-Triazole for Treatment of Non-Small Cell Lung Cancer. *Frontiers in Pharmacology*, 12. <https://doi.org/10.3389/fphar.2021.753676>
- [385] Yele, V., Azam, M. A., & Wadhwani, A. D. (2021). Synthesis, Molecular Docking and Biological Evaluation of 2-Aryloxy-N-Phenylacetamide and N'-(2-Aryloxyoxyacetyl) Benzohydrazide Derivatives as Potential Antibacterial Agents. *Chemistry and Biodiversity*, 18(4). <https://doi.org/10.1002/cbdv.202000907>
- [386] Zagórska, A., Bucki, A., Kołaczkowski, M., Siwek, A., Głuch-Lutwin, M., Starowicz, G., Kazek, G., Partyka, A., Wesołowska, A., Słoczyńska, K., Pękala, E., & Pawłowski, M. (2016). Synthesis and biological evaluation of 2-fluoro and 3-trifluoromethyl-phenyl-piperazinylalkyl derivatives of 1H-imidazo[2,1-f]purine-2,4(3H,8H)-dione as potential antidepressant agents. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 31, 10–24. <https://doi.org/10.1080/14756366.2016.1198902>
- [387] Zaheer, Z., Khan, F. A. K., Sangshetti, J. N., Patil, R. H., & Lohar, K. S. (2016). Novel amalgamation of phthalazine–quinolines as biofilm inhibitors: One-pot synthesis, biological evaluation and in silico ADME prediction with favorable metabolic fate. *Bioorganic and Medicinal Chemistry Letters*, 26(7), 1696–1703. <https://doi.org/10.1016/j.bmcl.2016.02.057>
- [388] Zhang, L., Chen, X., Xue, P., Sun, H. H. Y., Williams, I. D., Sharpless, K. B., Fokin, V. V., & Jia, G. (2005). Ruthenium-Catalyzed Cycloaddition of Alkynes and Organic Azides. *Journal of the American Chemical Society*, 127(46), 15998–15999. <https://doi.org/10.1021/ja054114s>

- [389] Zheng, A., Zhang, W., & Pan, J. (2006). One-Pot and Convenient Conversion of 5-Azidopyrazole-4-carboxaldehyde to Pyrazolo[3,4- *b*]pyridines. *Synthetic Communications*, 36(11), 1549–1556. <https://doi.org/10.1080/00397910600588868>
- [390] Zheng, X., Li, Z., Wang, Y., Chen, W., Huang, Q., Liu, C., & Song, G. (2003). Syntheses and insecticidal activities of novel 2,5-disubstituted 1,3,4-oxadiazoles. *Journal of Fluorine Chemistry*, 123(2), 163–169. [https://doi.org/10.1016/S0022-1139\(03\)00168-4](https://doi.org/10.1016/S0022-1139(03)00168-4)
- [391] Zhi, C., Long, Z., Manikowski, A., Comstock, J., Xu, W.-C., Brown, N. C., Tarantino, , Paul M., Holm, K. A., Dix, E. J., Wright, G. E., Barnes, M. H., Butler, M. M., Foster, K. A., LaMarr, W. A., Bachand, B., Bethell, R., Cadilhac, C., Charron, S., Lamothe, S., ... Storer, R. (2006). Hybrid Antibacterials. DNA Polymerase–Topoisomerase Inhibitors. *Journal of Medicinal Chemistry*, 49(4), 1455–1465. <https://doi.org/10.1021/jm0510023>
- [392] Zhu, D.-J., Chen, J.-X., Liu, M.-C., Ding, J.-C., & Wu, H.-Y. (2009). Catalyst: and solvent-free synthesis of imidazo[1,2-*a*]pyridines. *Journal of the Brazilian Chemical Society*, 20(3). <https://doi.org/10.1590/S0103-50532009000300012>
- [393] Zhu, Y. S., Xue, Y., Liu, W., Zhu, X., Hao, X. Q., & Song, M. P. (2020). Temperature-Controlled Chalcogenation and Chalcogenocyanation of Imidazopyridines in Water under Transition Metal-Free Conditions. *Journal of Organic Chemistry*, 85(14), 9106–9116. <https://doi.org/10.1021/acs.joc.0c01035>
- [394] Zhu, Z. Q., Guo, D., Ji, J. J., Zhu, X., Tang, J., Xie, Z. B., & Le, Z. G. (2020). Visible-Light-Induced Dehydrogenative Imidoylation of Imidazo[1,2- *a*]pyridines with α -Amino Acid Derivatives and α -Amino Ketones. *Journal of Organic Chemistry*, 85(23), 15062–15071. <https://doi.org/10.1021/acs.joc.0c01940>
- [395] Zisook, S., Mendels, J., Janowsky, D., Feighner, J., Lee, J. C. M., & Fritz, A. (1987). Efficacy and Safety of Fezolamine in Depressed Patients. *Neuropsychobiology*, 17(3), 133–138. <https://doi.org/10.1159/000118353>

Publications

International journals

- [1] Patel, M., & Kaneriya, D. (2024). Synthesis of 2-((5-2-methylimidazo[1,2-a]pyridine-3-yl)-1,3,4-oxadiazol-2-yl)thio)-N-phenylacetamide derivatives and their biological evaluation. *Rasayan journal of chemistry*.17(4).
<http://doi.org/10.31788/RJC.2024.1749004>

SCOPUS Indexed Journal-ISSN:0974-1496

- [2] Patel, M., & Kaneriya, D. (2024) Synthesis, Characterization, and Antimicrobial Activity of 1,2,3-Triazoles Containing a Theophylline Moiety. *Russian journal of organic chemistry*.60(12).
<http://doi.org/10.1134/S1070428024120182>

SCOPUS Indexed Journal-ISSN:1070-4280